antibodies .- online.com







anti-ARPC1A antibody (C-Term)





O	:
1 1\/\pi	view
\circ	V I C V V


Target:

Quantity:	400 μL
Target:	ARPC1A
Binding Specificity:	AA 286-315, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ARPC1A antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Flow Cytometry (FACS), Immunofluorescence (IF)
Product Details	
Immunogen:	This ARPC1A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 286-315 amino acids from the C-terminal region of human ARPC1A.
Clone:	RB17836
Isotype:	lg Fraction
Predicted Reactivity:	X, B, M, Rat
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Target Details	

ARPC1A


Target Details

Altornative Name:	ADDC1A (ADDC1A Producto)
Alternative Name:	ARPC1A (ARPC1A Products)
Background:	ARPC1A is one of seven subunits of the human Arp2/3 protein complex. This subunit is a
	member of the SOP2 family of proteins and is most similar to the protein ARPC1B. The
	similarity between these two proteins suggests that they both may function as p41 subunit of
	the human Arp2/3 complex that has been implicated in the control of actin polymerization in
	cells. It is possible that the p41 subunit is involved in assembling and maintaining the structure
	of the Arp2/3 complex. Multiple versions of the p41 subunit may adapt the functions of the
	complex to different cell types or developmental stages.
Molecular Weight:	41569
Gene ID:	10552
NCBI Accession:	NP_006400
UniProt:	Q92747
Pathways:	Regulation of Actin Filament Polymerization
Application Details	
Application Notes:	IF: 1:10~50. WB: 1:1000. IHC-P: 1:10~50. FC: 1:10~50
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in sma
	aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin-embedded human skeletal muscle reacted with ARPC1A Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated.



Western Blotting

Image 2. Anti-ARPC1A Antibody (C-term) at 1:1000 dilution + Hela whole cell lysate Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 42 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.



Immunofluorescence

Image 3. Immunofluorescence analysis of ARPC1A Antibody (C-term) with hela cells. 0.025 mg/mL primary antibody was followed by FITC-conjugated goat anti-rabbit IgG (whole molecule). FITC emits green fluorescence.Red counterstaining is PI.

Please check the product details page for more images. Overall 4 images are available for ABIN390417.