

Datasheet for ABIN390857
anti-SOD3 antibody (N-Term)

3 Images

1 Publication

[Go to Product page](#)

Overview

Quantity:	400 µL
Target:	SOD3
Binding Specificity:	AA 7-36, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SOD3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Flow Cytometry (FACS)

Product Details

Immunogen:	This SOD3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 7-36 amino acids from the N-terminal region of human SOD3.
Clone:	RB19866
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	SOD3
Alternative Name:	SOD3 (SOD3 Products)

Target Details

Background: SOD3 is a member of the superoxide dismutase(SOD) protein family. SODs are antioxidant enzymes that catalyze the dismutation of two superoxide radicals into hydrogen peroxide and oxygen. This protein is thought to protect the brain, lungs, and other tissues from oxidative stress. The protein is secreted into the extracellular space and forms a glycosylated homotetramer that is anchored to the extracellular matrix (ECM) and cell surfaces through an interaction with heparan sulfate proteoglycan and collagen. A fraction of the protein is cleaved near the C-terminus before secretion to generate circulating tetramers that do not interact with the ECM.

Molecular Weight: 25851

Gene ID: 6649

NCBI Accession: [NP_003093](#)

UniProt: [P08294](#)

Application Details

Application Notes: WB: 1:1000. IHC-P: 1:10~50. FC: 1:10~50

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Storage Comment: Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.

Expiry Date: 6 months

Publications


Product cited in: Murakami, Ito, Hagiwara, Yoshida, Sobue, Ichihara, Takagi, Kojima, Tanaka, Tamiya-Koizumi, Kyogashima, Suzuki, Banno, Nozawa, Murate: "ATRA inhibits ceramide kinase transcription in a

human neuroblastoma cell line, SH-SY5Y cells: the role of COUP-TFI." in: **Journal of neurochemistry**, Vol. 112, Issue 2, pp. 511-20, (2010) ([PubMed](#)).

Yang, Gagarin, St Laurent, Hammell, Toma, Hu, Iwasa, McCaffrey: "Cardiovascular inflammation and lesion cell apoptosis: a novel connection via the interferon-inducible immunoproteasome." in: **Arteriosclerosis, thrombosis, and vascular biology**, Vol. 29, Issue 8, pp. 1213-9, (2009) ([PubMed](#)).


Hinkovska-Galcheva, Clark, VanWay, Huang, Hiraoka, Abe, Borofsky, Kunkel, Shanley, Shayman, Lanni, Petty, Boxer: "Ceramide kinase promotes Ca²⁺ signaling near IgG-opsonized targets and enhances phagolysosomal fusion in COS-1 cells." in: **Journal of lipid research**, Vol. 49, Issue 3, pp. 531-42, (2008) ([PubMed](#)).

Images




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin-embedded human lung carcinoma reacted with SOD3 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated.



Western Blotting

Image 2. Western blot analysis of SOD3 Antibody (N-term) (ABIN390857 and ABIN2841079) in K562 cell line lysates (35 µg/lane). SOD3 (arrow) was detected using the purified Pab.



Flow Cytometry

Image 3. Flow cytometric analysis of K562 cells using SOD3 Antibody (N-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.