



Datasheet for ABIN390994
anti-MSK2 antibody (C-Term)



[Go to Product page](#)

1 Image

1 Publication

Overview

Quantity:	400 µL
Target:	MSK2 (RPS6KA4)
Binding Specificity:	AA 322-354, C-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MSK2 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	This RSKB (MSK2) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 322-354 amino acids from the C-terminal region of human RSKB (MSK2).
Clone:	RB1020
Isotype:	Ig Fraction
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Target Details

Target:	MSK2 (RPS6KA4)
Alternative Name:	RSKB (MSK2) (RPS6KA4 Products)

Target Details

Background: Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the γ phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The AGC kinase group consists of 63 kinases including the cyclic nucleotide-regulated protein kinase (PKA & PKG) family, the diacylglycerol-activated/phospholipid-dependent protein kinase C (PKC) family, the related to PKA and PKC (RAC/Akt) protein kinase family, the kinases that phosphorylate G protein-coupled receptors family (ARK), and the kinases that phosphorylate ribosomal protein S6 family (RSK). The calcium/calmodulin-dependent kinase (CAMK) group consists of 75 kinases regulated by $\text{Ca}^{2+}/\text{CaM}$ and close relative family (CAMK, CAMKL, DAPK, MAPKAPK).

Molecular Weight: 85606

Gene ID: 8986

NCBI Accession: [NP_001006945](#), [NP_003933](#)

UniProt: [O75676](#)

Application Details

Application Notes: WB: 1:1000

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small

Handling


aliquots to prevent freeze-thaw cycles.

Expiry Date: 6 months

Publications

Product cited in: Ewing, Chu, Elisma, Li, Taylor, Climie, McBroom-Cerajewski, Robinson, OConnor, Li, Taylor, Dharsee, Ho, Heilbut, Moore, Zhang, Ornatsky, Bukhman, Ethier, Sheng, Vasilescu, Abu-Farha, Lambert, Duewel et al.: "Large-scale mapping of human protein-protein interactions by mass spectrometry. ..." in: **Molecular systems biology**, Vol. 3, pp. 89, (2007) ([PubMed](#)).

Images



Western Blotting

Image 1. The anti-MSK2 Pab (ABIN390994 and ABIN2837937) is used in Western blot to detect MSK2 in placenta tissue lysate. Indirect immunofluorescence analysis showed that MSK2 is predominantly localized in the nucleus of parental (10T1/2) and oncogene-transformed (Ciras-3) mouse fibroblasts. Cells were co-stained with DI to visualize nucleus compartment. Data is kindly provided by B. Drobic and Dr. J. Davie from University of Manitoba (Winnipeg, Canada).