antibodies -online.com





anti-PKC gamma antibody (C-Term)

2 Images



Go to Product page

()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢

Target:

Quantity:	400 μL
Target:	PKC gamma (PRKCG)
Binding Specificity:	AA 305-336, C-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PKC gamma antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This PKC gamma antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 305-336 amino acids from the C-terminal region of human PKC gamma.
Immunogen: Clone:	synthetic peptide between 305-336 amino acids from the C-terminal region of human PKC
	synthetic peptide between 305-336 amino acids from the C-terminal region of human PKC gamma.
Clone:	synthetic peptide between 305-336 amino acids from the C-terminal region of human PKC gamma. RB1007
Clone:	synthetic peptide between 305-336 amino acids from the C-terminal region of human PKC gamma. RB1007 Ig Fraction

PKC gamma (PRKCG)


Target Details

Alternative Name:	PKC gamma (PRKCG Products)		
Background:	Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be		
	activated by calcium and second messenger diacylglycerol. PKC family members		
	phosphorylate a wide variety of protein targets and are known to be involved in diverse cellula		
	signaling pathways. PKC also serve as major receptors for phorbol esters, a class of tumor		
	promoters. Each member of the PKC family has a specific expression profile and is believed to		
	play distinct roles in cells. PKC gamma is one of the PKC family members. This protein kinase		
	is expressed solely in the brain and spinal cord and its localization is restricted to neurons. It		
	has been demonstrated that several neuronal functions, including long term potentiation (LTP		
	and long term depression (LTD), specifically require this kinase. Knockout studies in mice also		
	suggest that this kinase may be involved in neuropathic pain development. Defects in this		
	protein have been associated with neurodegenerative disorder spinocerebellar ataxia-14		
	(SCA14).		
Molecular Weight:	78448		
Gene ID:	5582		
NCBI Accession:	NP_002730		
UniProt:	P05129		
Pathways:	WNT Signaling, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Thyroid Hormone		
	Synthesis, Myometrial Relaxation and Contraction, G-protein mediated Events, Positive		
	Regulation of Response to DNA Damage Stimulus, Interaction of EGFR with phospholipase C-		
	gamma, Thromboxane A2 Receptor Signaling, VEGF Signaling		
Application Details			
Application Notes:	WB: 1:1000. IHC-P: 1:10~50		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which		

Handling

	should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in sn aliquots to prevent freeze-thaw cycles.	
Expiry Date:	6 months	

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin-embedded human brain tissue reacted with PKC gamma antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated.

Western Blotting

Image 2. Western blot analysis of anti-PKC gamma C-term Pab (ABIN391009 and ABIN2841182) in mouse brain lysate. PKC gamma (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.