antibodies -online.com





anti-SRPK1 antibody (N-Term)

2 Images



Go to Product page


Overview


Quantity:	400 μL
Target:	SRPK1
Binding Specificity:	AA 1-30, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SRPK1 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	This SRPK1 antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 1-30 amino acids from the N-terminal region of human SRPK1.
Clone:	RB9782
Isotype:	Ig Fraction
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by
	dialysis against PBS.
Target Details	
Target:	SRPK1
Alternative Name:	SRPK1 (SRPK1 Products)

Target Details

ranger became		
Background:	This gene encodes a serine/arginine protein kinase specific for the SR (serine/arginine-rich domain) family of splicing factors. The protein localizes to the nucleus and the cytoplasm. It is thought to play a role in regulation of both constitutive and alternative splicing by regulating intracellular localization of splicing factors. A second alternatively spliced transcript variant for this gene has been described, but its full length nature has not been determined.	
Molecular Weight:	74325	
Gene ID:	6732	
NCBI Accession:	NP_003128	
UniProt:	Q96SB4	
Pathways:	Toll-Like Receptors Cascades	
Application Details		
Application Notes:	WB: 1:2000. WB: 1:2000	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months





Western Blotting

Image 1. Anti-SRPK1 Antibody (N-term)at 1:2000 dilution + K562 whole cell lysate Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 74 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.

Western Blotting

Image 2. Anti-SRPK1 Antibody (N-term)at 1:2000 dilution + K562 whole cell lysate Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 74 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.