


Datasheet for ABIN391036

anti-PDPK1 antibody (N-Term)





Go to Product page

| () | ve | V / | - | 1 A |
|--------|--------|------|---|-----|
| | \cup | 1 \/ | - | 1/1 |
| \sim | ' V C | 1 V | ı | v v |
| | | | | |

| Overview | | |
|----------------------|---|--|
| Quantity: | 400 μL | |
| Target: | PDPK1 | |
| Binding Specificity: | AA 1-30, N-Term | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This PDPK1 antibody is un-conjugated | |
| Application: | Western Blotting (WB) | |
| Product Details | | |
| Immunogen: | This PDK1 antibody is generated from rabbits immunized with a KLH conjugated synthetic | |
| | peptide between 1-30 amino acids from the N-terminal region of human PDK1. | |
| Clone: | RB03036 | |
| Isotype: | lg Fraction | |
| Purification: | This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by | |
| | dialysis against PBS. | |
| Target Details | | |
| Target: | PDPK1 | |
| Alternative Name: | PDK1 (PDPK1 Products) | |
| | | |

Target Details

Expiry Date:

| rarget betails | | | |
|---------------------|---|--|--|
| Background: | Pyruvate dehydrogenase (PDH) is a mitochondrial multienzyme complex that catalyzes the | | |
| | oxidative decarboxylation of pyruvate and is one of the major enzymes responsible for the | | |
| | regulation of homeostasis of carbohydrate fuels in mammals. The enzymatic activity is | | |
| | regulated by a phosphorylation/dephosphorylation cycle. Phosphorylation of PDH by a specifi | | |
| | pyruvate dehydrogenase kinase (PDK) results in inactivation. | | |
| Molecular Weight: | 49244 | | |
| Gene ID: | 5163 | | |
| NCBI Accession: | NP_002601 | | |
| UniProt: | Q15118 | | |
| Pathways: | PI3K-Akt Signaling, TCR Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling | | |
| | Pathway, Neurotrophin Signaling Pathway, Regulation of Leukocyte Mediated Immunity, | | |
| | Positive Regulation of Immune Effector Process, Cell-Cell Junction Organization, Regulation o | | |
| | Cell Size, Skeletal Muscle Fiber Development, CXCR4-mediated Signaling Events, Signaling | | |
| | Events mediated by VEGFR1 and VEGFR2, VEGFR1 Specific Signals | | |
| Application Details | | | |
| Application Notes: | WB: 1:1000 | | |
| Restrictions: | For Research Use only | | |
| Handling | | | |
| Format: | Liquid | | |
| Buffer: | Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide. | | |
| Preservative: | Sodium azide | | |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which | | |
| | should be handled by trained staff only. | | |
| Storage: | 4 °C,-20 °C | | |
| Storage Comment: | Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in sm | | |
| | | | |


aliquots to prevent freeze-thaw cycles.

6 months

Product cited in:

Xu, Han, Epstein, Liu: "Regulation of PDK mRNA by high fatty acid and glucose in pancreatic islets." in: **Biochemical and biophysical research communications**, Vol. 344, Issue 3, pp. 827-33, (2006) (PubMed).

Images



Western Blotting

Image 1. Western blot analysis of PDK1 (arrow) using rabbit polyclonal hPDK1-G14 (ABIN391036 and ABIN2841202). 293 cell lysates (2 μg/lane) either nontransfected (Lane 1) or transiently transfected with the PDK1 gene (Lane 2) (Origene Technologies).