antibodies -online.com





anti-FGFR2 antibody (C-Term)

2 Images



Go to Product page

Overview

Overview	
Quantity:	400 μL
Target:	FGFR2
Binding Specificity:	AA 794-821, C-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FGFR2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This FGFR2 antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 794-821 amino acids from the C-terminal region of human FGFR2.
Clone:	RB1569
Isotype:	lg Fraction
Predicted Reactivity:	М
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	FGFR2
Alternative Name:	FGFR2 (FGFR2 Products)


FGFR2 is a member of the fibroblast growth factor receptor family, where amino acid sequence
is highly conserved between members and throughout evolution. FGFR family members differ
from one another in their ligand affinities and tissue distribution. A full-length representative
protein consists of an extracellular region, composed of three immunoglobulin-like domains, a
single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain.
The extracellular portion of the protein interacts with fibroblast growth factors, setting in
motion a cascade of downstream signals, ultimately influencing mitogenesis and
differentiation. This particular family member is a high-affinity receptor for acidic, basic and/or
keratinocyte growth factor, depending on the isoform. Mutations in the gene are associated
with many craniosynostotic syndromes and bone malformations. The genomic organization of
the gene encompasses 20 exons. Alternative splicing in multiple exons, including those
encoding the Ig-like domains, the transmembrane region and the carboxyl terminus, results in
varied isoforms which differ in structure and specificity. Isoform 1 has equal affinity for aFGF and bFGF but does not bind KGF.
and bror but does not bind Nor.
92025
2263
NP_000132, NP_001138385, NP_001138386, NP_001138387, NP_001138388, NP_001138389,
NP_001138390, NP_001138391, NP_075259, NP_075418
P21802
RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin
Signaling Pathway, Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber
Development, Growth Factor Binding


Application Notes:	WB: 1:1000. IHC-P-Leica: 1:500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

Handling

	should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Images





Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Immunohistochemical analysis of paraffinembedded Human breast carcinoma tissue using B performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH 9. 0). Samples were incubated with primary Antibody (1:500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.

Western Blotting

Image 2. All lanes: Anti-FGFR2 Antibody (C-term) at 1:1000 dilution Lane 1: NCI- whole cell lysate Lane 2: HT-29 whole cell lysate Lane 3: A549 whole cell lysate Lane 4: T47D whole cell lysate Lane 5: Hela whole cell lysate Lane 6: NIH/3T3 whole cell lysate Lane 7: Mouse brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 92 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.