

Datasheet for ABIN392230  
**anti-PTK7 antibody (N-Term)**



[Go to Product page](#)

3 Images 1 Publication

## Overview

Quantity:	400 µL
Target:	PTK7
Binding Specificity:	AA 21-52, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PTK7 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Immunogen:	This CCK4 (PTK7) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 21-52 amino acids from the N-terminal region of human CCK4 (PTK7).
Clone:	RB01501
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

## Target Details

Target:	PTK7
Alternative Name:	CCK4 (PTK7) ( <a href="#">PTK7 Products</a> )

## Target Details

Background:	CCK4 may function as a cell adhesion molecule. Although it belongs to the insuline receptor subfamily of the Tyr protein kinases, it likely lacks the catalytic activity of a tyrosine kinase. It may be connected to the pathophysiology of colon carcinomas and/or may represent a tumor progression marker. This Type I membrane protein is highly expressed in lung, liver, pancreas, kidney, placenta and melanocytes, but weakly expressed in thyroid gland, ovary, brain, heart and skeletal muscle, and not in colon. It is also expressed in erythroleukemia cells.
Molecular Weight:	118392
Gene ID:	5754
NCBI Accession:	<a href="#">NP_001257327</a> , <a href="#">NP_002812</a> , <a href="#">NP_690619</a> , <a href="#">NP_690620</a> , <a href="#">NP_690621</a>
UniProt:	<a href="#">Q13308</a>
Pathways:	<a href="#">RTK Signaling</a> , <a href="#">Tube Formation</a>

## Application Details

Application Notes:	WB: 1:1000-1:2000. IHC-P: 1:50~100. FC: 1:10~50
Restrictions:	For Research Use only




## Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

## Publications

Product cited in:	Wang, Li, Yin, Wang, Wang, Wei, Guo, Ma, Shi, Zhou, Wang: "PTK7 protein is decreased in epithelial ovarian carcinomas with poor prognosis." in: <b>International journal of clinical and</b>
-------------------	--

Images



Flow Cytometry

**Image 1.** Flow cytometric analysis of NCI- cells using CCK4 (PTK7) Antibody (N-term) (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Western Blotting

**Image 2.** All lanes : Anti-CCK4 (PTK7) Antibody (N-term) at 1:1000-1:2000 dilution Lane 1: A431 whole cell lysate Lane 2: HeLa whole cell lysate Lane 3: HepG2 whole cell lysate Lane 4: Caco2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 118 kDa Blocking/Dilution buffer: 5 % NFDm/TBST.

Immunohistochemistry (Paraffin-embedded Sections)

**Image 3.** Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated. BC = breast carcinoma, HC = hepatocarcinoma.