antibodies -online.com





anti-TRAF2 antibody (C-Term)

2 Images



Go to Product page

)\/(

Overview	
Quantity:	400 μL
Target:	TRAF2
Binding Specificity:	AA 343-372, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRAF2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This TRAF2 antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 343-372 amino acids from the C-terminal region of human TRAF2.
Clone:	RB13076
Isotype:	Ig Fraction
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by
	dialysis against PBS.
Target Details	
Target:	TRAF2
Alternative Name:	TRAF2 (TRAF2 Products)

Target Details

Background:

TRAF2 is a member of the TNF receptor associated factor (TRAF) protein family. TRAF proteins associate with, and mediate the signal transduction from members of the TNF receptor superfamily. This protein directly interacts with TNF receptors, and forms a heterodimeric complex with TRAF1. This protein is required for TNF-alpha-mediated activation of MAPK8/JNK and NF-kappaB. The protein complex formed by this protein and TRAF1 interacts with the inhibitor-of-apoptosis proteins (IAPs), and functions as a mediator of the anti-apoptotic signals from TNF receptors. The interaction of this protein with TRADD, a TNF receptor associated apoptotic signal transducer, ensures the recruitment of IAPs for the direct inhibition of caspase activation. BIRC2/c-IAP1, an apoptosis inhibitor possessing ubiquitin ligase activity, can unbiquitinate and induce the degradation of this protein, and thus potentiate TNF-induced apoptosis.

Molecular Weight:	55859
Gene ID:	7186
NCBI Accession:	NP_066961
UniProt:	Q12933
Pathways:	NF-kappaB Signaling, Apoptosis, Caspase Cascade in Apoptosis, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, Positive Regulation of Endopeptidase Activity, Hepatitis C,

Unfolded Protein Response, S100 Proteins

should be handled by trained staff only.

4 °C,-20 °C

Application Details

Precaution of Use:

Storage:

Application Notes:	WB: 1:1000. IHC-P: 1:10~50
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide

This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which


Handling


Storage Comment: Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small

aliquots to prevent freeze-thaw cycles.

Expiry Date: 6 months

Images





Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin-embedded human colon carcinoma tissue reacted with TRAF2 antibody (Cterm) (ABIN392282 and ABIN2841948), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated.

Western Blotting

Image 2. Western blot analysis of TRAF2 (arrow) using rabbit polyclonal TRAF2 Antibody (C-term) (ABIN392282 and ABIN2841948). 293 cell lysates ($2 \mu g$ /lane) either nontransfected (c) or transiently transfected with the TRAF2 gene (Lane 2) (Origene Technologies).