antibodies .- online.com





anti-Transglutaminase 2 antibody (AA 429-458)





Go to Product page

\sim	
()\/\	rview
\circ	


Target:

Overview	
Quantity:	400 μL
Target:	Transglutaminase 2 (TGM2)
Binding Specificity:	AA 429-458
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Transglutaminase 2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This Transglutaminase (TGM2) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 429-458 amino acids from the Central region of human Transglutaminase (TGM2).
Clone:	RB13081
Isotype:	lg Fraction
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Target Details	
_	

Transglutaminase 2 (TGM2)


Target Details

rarget Details	
Alternative Name:	Transglutaminase (TGM2) (TGM2 Products)
Background:	Transglutaminases are enzymes that catalyze the crosslinking of proteins by epsilon-gamma glutamyl lysine isopeptide bonds. While the primary structure of transglutaminases is not conserved, they all have the same amino acid sequence at their active sites and their activity is calcium-dependent. TGM2 acts as a monomer, is induced by retinoic acid, and appears to be involved in apoptosis. It has been identified as the autoantigen implicated in celiac disease.
Molecular Weight:	77329
Gene ID:	7052
NCBI Accession:	NP_004604, NP_945189
UniProt:	P21980
Pathways:	Tube Formation, Thromboxane A2 Receptor Signaling
Application Details	
Application Notes:	IF: 1:10~50. WB: 1:1000. WB: 1:1000. IHC-P: 1:10~50
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with *TGM2 antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated.



Western Blotting

Image 2. TGM2 Antibody (Center) (ABIN392285 and ABIN2841950) western blot analysis in HUVEC,K562 cell line lysates (35 μ g/lane).This demonstrates the TGM2 antibody detected the TGM2 protein (arrow).



Immunofluorescence

Image 3. Fluorescent confocal image of cell stained with TGM2 Antibody (Center) (ABIN392285 and ABIN2841950). cells were fixed with 4 % PFA (20 min), permeabilized with Triton X-100 (0.1 %, 10 min), then incubated with TGM2 primary antibody (1:25, 1 h at 37 °C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit (1:400,50 min antibody (green) was used 37 °C). Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7 units/mL, 1 h at 37 °C). Nuclei were counterstained with DI (blue) (10 μg/mL, 10 min).TGM2 immunoreactivity is localized to Cytoplasm significantly.

Please check the product details page for more images. Overall 4 images are available for ABIN392285.