antibodies -online.com







anti-PLK1 antibody (C-Term)

Images

Publications



| \sim | |
|--------|----------|
| ()\/\ | rview |
| \cup | 1 410 44 |

| 0.00000 | |
|-----------------------|---|
| Quantity: | 400 μL |
| Target: | PLK1 |
| Binding Specificity: | AA 573-603, C-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Application: | Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |
| Product Details | |
| Immunogen: | This PLK1 antibody is generated from rabbits immunized with a KLH conjugated synthetic |
| | peptide between 573-603 amino acids from the C-terminal region of human PLK1. |
| Clone: | RB00878 |
| Isotype: | Ig Fraction |
| Predicted Reactivity: | М |
| Purification: | This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by |
| | dialysis against PBS. |
| Target Details | |
| Target: | PLK1 |
| Alternative Name: | PLK1 (PLK1 Products) |

Target Details

| Background: |
|-------------|
|-------------|

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the g phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains.

| Molecular Weight: | 68255 |
|-------------------|------------------------------|
| Gene ID: | 5347 |
| NCBI Accession: | NP_005021 |
| UniProt: | P53350 |
| Pathways: | Cell Division Cycle, M Phase |

Application Details

Restrictions: For Research Use only


Handling

| Format: | Liquid |
|--------------------|--|
| Buffer: | Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles. |
| Expiry Date: | 6 months |

Product cited in:

Sadeghi, Ullenhag, Wagenius, Tötterman, Eriksson: "Rapid expansion of T cells: Effects of culture and cryopreservation and importance of short-term cell recovery." in: **Acta oncologica** (Stockholm, Sweden), Vol. 52, Issue 5, pp. 978-86, (2013) (PubMed).

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated. BC = breast carcinoma, HC = hepatocarcinoma.

Western Blotting

Image 2. Western blot analysis of anti-PLK Pab (ABIN392446 and ABIN2842043) in cell lysate. PLK (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.