

Datasheet for ABIN392462
anti-SGK3 antibody (N-Term)



[Go to Product page](#)

4 Images

2 Publications

Overview

Quantity:	400 µL
Target:	SGK3
Binding Specificity:	AA 1-30, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SGK3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This SGK3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human SGK3.
Clone:	RB0888
Isotype:	Ig Fraction
Predicted Reactivity:	M, Rat
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Target Details

Target:	SGK3
---------	------

Target Details

Alternative Name:	SGK3 (SGK3 Products)
Background:	SGK3, a Ser/Thr protein kinase, is similar to serum- and glucocorticoid-induced protein kinase (SGK), but this gene product is not induced by serum or glucocorticoids. Expression is induced in response to signals that activate phosphatidylinositol 3-kinase, which is also true for SGK.
Molecular Weight:	57108
Gene ID:	23678
NCBI Accession:	NP_001028750 , NP_001191102 , NP_037389 , NP_733827
UniProt:	Q96BR1

Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:50~100. IHC-P: 1:50~100. IHC-P: 1:50~100
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months




Publications

Product cited in: Vadnais, Aghajanian, Lin, Gerton: "Signaling in sperm: toward a molecular understanding of the acquisition of sperm motility in the mouse epididymis." in: **Biology of reproduction**, Vol. 89, Issue 5, pp. 127, (2013) ([PubMed](#)).

Okada, Ishii, Masujin, Yasoshima, Matsuda, Ogura, Nakayama, Kunieda, Doi: "The critical roles

of serum/glucocorticoid-regulated kinase 3 (SGK3) in the hair follicle morphogenesis and homeostasis: the allelic difference provides novel insights into hair follicle biology." in: **The American journal of pathology**, Vol. 168, Issue 4, pp. 1119-33, (2006) ([PubMed](#)).

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. IHC detection of SGK3 protein on the paraffin sections of the WT (left) and YPC (right) mice at P0 (B and C), P3 (D and E), and P5 (F and G) skin. Positive signals were observed in the cytoplasm of the hair follicle keratinocytes, especially in hair bulb, ORS, IRS, cuticle/cortex and bulge, or sebaceous glands. Some differences between the WT and YPC, for example, the expression in bulb keratinocytes were observed at P3 and P5. Scale bar, 50.

Western Blotting

Image 2. Western blot analysis of anti-SGK3 Pab (ABIN392462 and ABIN2842050) in cell lysate. SGK3 (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.

Immunohistochemistry (Paraffin-embedded Sections)

Image 3. Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated. BC = breast carcinoma, HC = hepatocarcinoma.

Images

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN392462.