antibodies -online.com





anti-PACSIN1 antibody (N-Term)



Images



Go to Product page

()	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	$I \vee I$	ew

Overview	
Quantity:	400 μL
Target:	PACSIN1
Binding Specificity:	AA 8-38, N-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PACSIN1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This PACSIN1 antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 8-38 amino acids from the N-terminal region of human PACSIN1.
Clone:	RB3586
Isotype:	lg Fraction
Predicted Reactivity:	B, C, X
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	PACSIN1
Alternative Name:	PACSIN1 (PACSIN1 Products)

Target Details

Background:

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the g phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The STE group (homologs of yeast Sterile 7, 11, 20 kinases) consists of 50 kinases related to the mitogen-activated protein kinase (MAPK) cascade families (Ste7/MAP2K, Ste11/MAP3K, and Ste20/MAP4K). MAP kinase cascades, consisting of a MAPK and one or more upstream regulatory kinases (MAPKKs) have been best characterized in the yeast pheromone response pathway. Pheromones bind to Ste cell surface receptors and activate yeast MAPK pathway.

Molecular Weight:	50966
Gene ID:	29993
NCBI Accession:	NP_001186512, NP_065855
UniProt:	Q9BY11

Application Details

Application Notes:	WB: 1:1000.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.

Expiry Date:


6 months

Images




Western Blotting

Image 1. Western blot analysis of CSIN1 (arrow) using rabbit polyclonal CSIN1 Antibody (G23) (ABIN392664 and ABIN2842162). 293 cell lysates (2 μ g/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the CSIN1 gene.



Western Blotting

Image 2. Western blot analysis of lysates from Daudi cell line, human brain, human ncrease, human heart tissue (from left to right), using CSIN1 Antibody (G23) (ABIN392664 and ABIN2842162). (ABIN392664 and ABIN2842162) was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20 µg per lane.



Western Blotting

Image 3. Western blot analysis of lysates from Daudi cell line, human heart and ncreas tissue lysate(from left to right), using CSIN1 Antibody (G23) (ABIN392664 and ABIN2842162). (ABIN392664 and ABIN2842162) was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 35 μg per lane.

Please check the product details page for more images. Overall 7 images are available for ABIN392664.