antibodies -online.com





anti-PFKFB3 antibody (C-Term)

3 Images

Target:



PFKFB3

Publications



Go to Product page

Overview	
Quantity:	400 μL
Target:	PFKFB3
Binding Specificity:	AA 454-484, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PFKFB3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This PFKFB3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 454-484 amino acids from the C-terminal region of human PFKFB3.
Clone:	RB04030
Isotype:	lg Fraction
Predicted Reactivity:	Rat
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Target Details	

Target Details

Alternative Name:	PFKFB3 (PFKFB3 Products)
Background:	Protein kinases are enzymes that transfer a phosphate group from a phosphate donor,
	generally the g phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this
	basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells,
	regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement
	and cell movement, apoptosis, and differentiation. With more than 500 gene products, the
	protein kinase family is one of the largest families of proteins in eukaryotes. The family has
	been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or
	serine/threonine (STK) kinase catalytic domains. The AGC kinase group consists of 63 kinases
	including the cyclic nucleotide-regulated protein kinase (PKA & PKG) family, the diacylglycerol-
	activated/phospholipid-dependent protein kinase C (PKC) family, the related to PKA and PKC
	(RAC/Akt) protein kinase family, the kinases that phosphorylate G protein-coupled receptors
	family (ARK), and the kinases that phosphorylate ribosomal protein S6 family (RSK).
Molecular Weight:	59609
Gene ID:	5209
NCBI Accession:	NP_001138915, NP_004557
UniProt:	Q16875
Pathways:	AMPK Signaling, Regulation of Carbohydrate Metabolic Process
Application Details	
Application Notes:	WB: 1:1000. WB: 1:1000. IHC-P: 1:50~100
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Handling Advice:	Avoid freeze-thaw cycles.
Storage:	4 °C,-20 °C

Handling

Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small Storage Comment: aliquots.

6 months

Expiry Date:

Publications

Product cited in:

Heydasch, Kessler, Warnke, Eschrich, Scholz, Bigl: "Functional diversity of PFKFB3 splice variants in glioblastomas." in: PloS one, Vol. 16, Issue 7, pp. e0241092, (2021) (PubMed).


Lee, Lee, Yun, Jang, Kang, Kim, Choi, Park: "Silver nanoparticles affect glucose metabolism in hepatoma cells through production of reactive oxygen species." in: International journal of nanomedicine, Vol. 11, pp. 55-68, (2016) (PubMed).


Reddy, Fernandes, Deshpande, Weisberg, Inguilizian, Abdel-Wahab, Kung, Levine, Griffin, Sattler: "The JAK2V617F oncogene requires expression of inducible phosphofructokinase/fructosebisphosphatase 3 for cell growth and increased metabolic activity." in: Leukemia, Vol. 26, Issue 3, pp. 481-9, (2012) (PubMed).

Ando, Uehara, Kogure, Asano, Nakajima, Abe, Kawauchi, Tanaka: "Interleukin 6 enhances glycolysis through expression of the glycolytic enzymes hexokinase 2 and 6-phosphofructo-2kinase/fructose-2,6-bisphosphatase-3." in: Journal of Nippon Medical School = Nippon Ika Daigaku zasshi, Vol. 77, Issue 2, pp. 97-105, (2010) (PubMed).

Yamasaki, Hayashi, Okamoto, Osanai, Lee: "Insulin-independent promotion of chemically induced hepatocellular tumor development in genetically diabetic mice." in: Cancer science, Vol. 101, Issue 1, pp. 65-72, (2010) (PubMed).

There are more publications referencing this product on: Product page





Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated. BC = breast carcinoma, HC = hepatocarcinoma.

Western Blotting

Image 2. Western blot analysis of lysate from human kidney tissue lysate, using PFKFB3 Antibody (ABIN392768 and ABIN2842213). (ABIN392768 and ABIN2842213) was diluted at 1:1000. A goat anti-rabbit lgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35 μ g.

Western Blotting

Image 3. Western blot analysis of anti-PFKFB3 Antibody (Cterm) (ABIN392768 and ABIN2842213) in CEM cell line lysates (35 μ g/lane). PFKFB3(arrow) was detected using the purified Pab.