.-online.com antibodies

Datasheet for ABIN411278 HGF ELISA Kit

1 Image

4 Publications



Overview

Quantity:	96 tests
Target:	HGF
Binding Specificity:	AA 32-494, AA 495-728
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	125-8000 pg/mL
Minimum Detection Limit:	125 pg/mL
Application:	ELISA

Product Details

Purpose:	Sandwich High Sensitivity ELISA kit for Quantitative Detection of Human HGF
Brand:	PicoKine™
Sample Type:	Cell Culture Supernatant, Serum, Plasma (EDTA), Plasma (citrate)
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Immunogen:	Expression system for standard: sf21 Immunogen sequence: Q32-R494(alpha)&V495-S728(beta)
Specificity:	Expression system for standard: sf21 Immunogen sequence: Q32-R494(alpha)&V495-S728(beta)
Cross-Reactivity (Details):	There is no detectable cross-reactivity with other relevant proteins.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN411278 | 11/30/2023 | Copyright antibodies-online. All rights reserved.

Product Details

Sensitivity:	<5pg/mL
Material not included:	Microplate reader in standard size. Automated plate washer. Adjustable pipettes and pipette
	tips. Multichannel pipettes are recommended in the condition of large amount of samples in the
	detection. Clean tubes and Eppendorf tubes. Washing buffer (neutral PBS or TBS). Preparation
	of 0.01M TBS: Add 1.2g Tris, 8.5g Nacl

Target Details

Target:	HGF
Alternative Name:	HGF (HGF Products)
Background:	Protein Function: Potent mitogen for mature parenchymal hepatocyte cells, seems to be a
	hepatotrophic factor, and acts as a growth factor for a broad spectrum of tissues and cell
	types. Activating ligand for the receptor tyrosine kinase MET by binding to it and promoting its
	dimerization
	Background: Hepatocyte growth factor(HGF) is the most potent mitogen for mature
	parenchymal hepatocytes in primary culture, and seems to be a hepatotrophic factor that acts
	as a trigger for liver regeneration after partial hepatectomy and liver injury. HGF has a relative
	molecular mass(Mr) of 82,000 and is a heterodimer composed of a large alpha-subunit of Mr
	69,000 and a small beta-subunit of Mr 34,000. The protein consists of 728 amino acid residues,
	including a possible signal peptide at the N-terminus. HGF may serve as a paracrine mediator
	to control placental development and growth. This growth factor may play an important role as
	a paracrine mediator of the proliferation of melanocytes and endothelial cells, as well as cells of
	epithelial origin. The gene encoding the human HGF is assigned to human chromosome 7.
	Synonyms: Hepatocyte growth factor,Hepatopoietin-A,Scatter factor,SF,Hepatocyte growth
	factor alpha chain,Hepatocyte growth factor beta chain,HGF,HPTA,
	Full Gene Name: Hepatocyte growth factor
Gene ID:	3082
UniProt:	P14210
Pathways:	RTK Signaling, Carbohydrate Homeostasis, Glycosaminoglycan Metabolic Process, Synaptic
	Membrane, Signaling of Hepatocyte Growth Factor Receptor

Application Details

Application Notes:	Before using Kit, spin tubes and bring down all components to bottom of tube. Duplicate well
	assay was recommended for both standard and sample testing.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN411278 | 11/30/2023 | Copyright antibodies-online. All rights reserved.

· · ·	
Comment:	Sequence similarities: Belongs to the peptidase S1 family. Plasminogen subfamily.
Plate:	Pre-coated
Protocol:	human HGF ELISA Kit was based on standard sandwich enzyme-linked immune-sorbent assay technology. A monoclonal antibody from mouse specific for HGF has been precoated onto 96- well plates. Standards(sf21, Q32-R494(alpha) & V495-S728(beta)) and test samples are added to the wells, a biotinylated detection polyclonal antibody from goat specific for HGF is added subsequently and then followed by washing with PBS or TBS buffer. Avidin-Biotin-Peroxidase Complex was added and unbound conjugates were washed away with PBS or TBS buffer. HRP substrate TMB was used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the human HGF amount of sample captured in plate.
Assay Procedure:	Aliquot 0.1 mL per well of the 8000pg/mL,4000pg/mL, 2000pg/mL, 1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL human HGF standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of human cell culture supernates, serum or plasma(EDTA, citrate) to each empty well. See "Sample Dilution Guideline" above for details. It is recommended that each human HGF standard solution and each sample be measured in duplicate.
Assay Precision:	 Sample 1: n=16, Mean(pg/ml): 521, Standard deviation: 39.6, CV(%): 7.6 Sample 2: n=16, Mean(pg/ml): 3649, Standard deviation: 222.6, CV(%): 6.1 Sample 3: n=16, Mean(pg/ml): 5621, Standard deviation: 269.8, CV(%): 4.8, Sample 1: n=24, Mean(pg/ml): 532, Standard deviation: 40.96, CV(%): 7.7 Sample 2: n=24, Mean(pg/ml): 3576, Standard deviation: 303.96, CV(%): 8.5 Sample 3: n=24, Mean(pg/ml): 5618, Standard deviation: 348.3, CV(%): 6.2
Restrictions:	For Research Use only
Handling	
Handling Advice:	Avoid multiple freeze-thaw cycles.
Storage:	-20 °C,4 °C
Storage Comment:	Store at 4°C for 6 months, at -20°C for 12 months. Avoid multiple freeze-thaw cycles
Expiry Date:	12 months
Product cited in: Order at www.a	Li, Hu, Wang, Zhang, Zhou, Yang, Li, Xiong, Liu, Li, Wu, Zheng: "Autophagy-dependent generation

Application Details

International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN411278 | 11/30/2023 | Copyright antibodies-online. All rights reserved. of Axin2+ cancer stem-like cells promotes hepatocarcinogenesis in liver cirrhosis." in: **Oncogene**, Vol. 36, Issue 48, pp. 6725-6737, (2017) (PubMed).

Maraldi, Beretti, Guida, Zavatti, De Pol: "Role of hepatocyte growth factor in the immunomodulation potential of amniotic fluid stem cells." in: **Stem cells translational medicine**, Vol. 4, Issue 6, pp. 539-47, (2015) (PubMed).

Guo, Sun, Xu, Zhao, Peng, Wang: "Human umbilical cord mesenchymal stem cells promote peripheral nerve repair via paracrine mechanisms." in: **Neural regeneration research**, Vol. 10, Issue 4, pp. 651-8, (2015) (PubMed).

Secchiero, Corallini, Zavan, Tripodo, Vindigni, Zauli: "Mesenchymal stem cells display hepatoprotective activity in lymphoma bearing xenografts." in: **Investigational new drugs**, Vol. 30, Issue 2, pp. 803-7, (2012) (PubMed).

Images



ELISA

Image 1. Human HGF PicoKine ELISA Kit standard curve