

# Datasheet for ABIN411283

# Interferon gamma ELISA Kit

1 Image

33

Publications



Go to Product page

### Overview

Quantity:	96 tests
Target:	Interferon gamma (IFNG)
Binding Specificity:	AA 23-155
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	31.2-2000 pg/mL
Minimum Detection Limit:	31.2 pg/mL
Application:	ELISA

## **Product Details**

Purpose:	Sandwich High Sensitivity ELISA kit for Quantitative Detection of Mouse IFN gamma
Brand:	PicoKine™
Sample Type:	Cell Culture Supernatant, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Immunogen:	Expression system for standard: E.coli Immunogen sequence: H23-C155
Specificity:	Expression system for standard: E.coli Immunogen sequence: H23-C155
Cross-Reactivity (Details):	There is no detectable cross-reactivity with other relevant proteins.

Product Details	
Sensitivity:	<5pg/mL
Material not included:	Microplate reader in standard size. Automated plate washer. Adjustable pipettes and pipette
	tips. Multichannel pipettes are recommended in the condition of large amount of samples in the
	detection. Clean tubes and Eppendorf tubes. Washing buffer (neutral PBS or TBS). Preparation
	of 0.01M TBS: Add 1.2g Tris, 8.5g Nacl
Target Details	
Target:	Interferon gamma (IFNG)
Alternative Name:	IFNG (IFNG Products)
Background:	Protein Function: Produced by lymphocytes activated by specific antigens or mitogens. IFN-
	gamma, in addition to having antiviral activity, has important immunoregulatory functions. It is
	a potent activator of macrophages, it has antiproliferative effects on transformed cells and it
	can potentiate the antiviral and antitumor effects of the type I interferons.
	Background: Interferon-gamma(IFN-gamma) is an inflammatory cytokine that has been
	implicated in the development of fibrosis in inflamed tissues. The production of IFN-gamma,
	which is under genetic control, can influence the development of fibrosis in lung allografts. IFN-
	gamma is also produced by natural killer(NK) cells and most prominently by CD8 cytotoxic T
	cells, and is vital for the control of microbial pathogens. Interferon gamma is believed to be
	crucial for host defence against many infections. Genetically determined variability in IFN-
	gamma and expression might be important for the development of tuberculosis. IFN-gamma
	activates human macrophage oxidative metabolism and antimicrobial activity. In addition to
	having antiviral activity, IFN-gamma has important immunoregulatory functions. IFN-gamma

Synonyms: Interferon gamma, IFN-gamma, Ifng,

plays an important role in the control of neointima proliferation.

Full Gene Name: Interferon gamma

Cellular Localisation: Secreted.

Gene ID: 15978

UniProt: P01580

Pathways:

Interferon-gamma Pathway, Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, ER-Nucleus Signaling, Regulation of Carbohydrate Metabolic Process, Protein targeting to Nucleus, Autophagy

# **Application Details**

Application Notes:	Before using Kit, spin tubes and bring down all components to bottom of tube. Duplicate well assay was recommended for both standard and sample testing.
Comment:	Tissue Specificity: Released primarily from activated T lymphocytes.
Plate:	Pre-coated
Protocol:	mouse IFN gamma ELISA Kit was based on standard sandwich enzyme-linked immune-sorbent
	assay technology. A monoclonal antibody from rat specific for IFN gamma has been precoated
	onto 96-well plates. Standards(E.coli, H23-C155) and test samples are added to the wells, a
	biotinylated detection polyclonal antibody from goat specific for IFN gamma is added
	subsequently and then followed by washing with PBS or TBS buffer. Avidin-Biotin-Peroxidase
	Complex was added and unbound conjugates were washed away with PBS or TBS buffer. HRP
	substrate TMB was used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to
	produce a blue color product that changed into yellow after adding acidic stop solution. The
	density of yellow is proportional to the mouse IFN gamma amount of sample captured in plate.
Assay Procedure:	Aliquot 0.1 mL per well of the 2000pg/mL,1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL,
	62.5pg/mL, 31.2pg/mL mouse IFN gamma standard solutions into the precoated 96-well plate.
	Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each
	properly diluted sample of mouse cell culture supernates or serum to each empty well. See
	"Sample Dilution Guideline" above for details. It is recommended that each mouse IFN gamma
	standard solution and each sample be measured in duplicate.
Assay Precision:	• Sample 1: n=16, Mean(pg/ml): 305, Standard deviation: 11.59, CV(%): 3.8
	• Sample 2: n=16, Mean(pg/ml): 782, Standard deviation: 35.19, CV(%): 4.5
	<ul> <li>Sample 3: n=16, Mean(pg/ml): 1367, Standard deviation: 69.72, CV(%): 5.1,</li> <li>Sample 1: n=24, Mean(pg/ml): 350, Standard deviation: 19.6, CV(%): 5.6</li> </ul>
	• Sample 2: n=24, Mean(pg/ml): 812, Standard deviation: 15.0, CV(%): 6.8
	Sample 3: n=24, Mean(pg/ml): 1494, Standard deviation: 109.1, CV(%): 7.3
Restrictions:	For Research Use only
Handling	
Handling Advice:	Avoid multiple freeze-thaw cycles.
Storage:	-20 °C,4 °C
Storage Comment:	Store at 4°C for 6 months, at -20°C for 12 months. Avoid multiple freeze-thaw cycles
Expiry Date:	12 months

Product cited in:

Qin, Gao, Liu, Liu, Wang, Zou: "Silencing of suppressor of cytokine signaling 1 enhances the immunological effect of mucin 1-calreticulin-primed 4T1 cell-treated dendritic cells in breast cancer treatment." in: **Oncology letters**, Vol. 15, Issue 2, pp. 1630-1638, (2018) (PubMed).

Lu, Liu, Jin, Chen, Liang, Qiu, Dai: "Herbal Components of a Novel Formula PSORI-CM02 Interdependently Suppress Allograft Rejection and Induce CD8+CD122+PD-1+ Regulatory T Cells." in: **Frontiers in pharmacology**, Vol. 9, pp. 88, (2018) (PubMed).


Hu, Feng, Wang, Song, Ni: "5-BDBD ameliorates an OVA-induced allergic asthma by the reduction of Th2 cytokines production." in: **Iranian journal of basic medical sciences**, Vol. 21, Issue 4, pp. 364-369, (2018) (PubMed).

Zhang, Li, Wang, Xie, Li, Zuo, Kong, Wang, Wang: "Immune Protection of Rhoptry Protein 21 (ROP21) of Toxoplasma gondii as a DNA Vaccine Against Toxoplasmosis." in: **Frontiers in microbiology**, Vol. 9, pp. 909, (2018) (PubMed).

Xu, Liu, Wang, Li, Wu, Li, Di, You, Jiang et al.: "3-(1H-Benzo[d]imidazol-6-yl)-5-(4-fluorophenyl)-1,2,4-oxadiazole (DD07232), a Novel Potent Nrf2/ARE Inducer, Ameliorates DSS-Induced Murine Colitis and Protects NCM460 Cells against Oxidative Stress ..." in: **Oxidative medicine and cellular longevity**, Vol. 2018, pp. 3271617, (2018) (PubMed).

There are more publications referencing this product on: Product page

### **Images**



### **ELISA**

Image 1. Mouse IFN gamma PicoKine ELISA Kit standard curve