



[Go to Product page](#)

Datasheet for ABIN411292

IL1A ELISA Kit

1 Image

13 Publications

Overview

Quantity: 96 tests

Target: IL1A

Binding Specificity: AA 115-270

Reactivity: Rat

Method Type: Sandwich ELISA

Detection Range: 4.69-300 pg/mL

Minimum Detection Limit: 4.69 pg/mL

Application: ELISA

Product Details

Purpose: Sandwich High Sensitivity ELISA kit for Quantitative Detection of Rat IL-1 alpha

Brand: PicoKine™

Sample Type: Cell Culture Supernatant, Serum

Analytical Method: Quantitative

Detection Method: Colorimetric

Immunogen: Expression system for standard: E.coli
Immunogen sequence: S115-S270

Specificity: Expression system for standard: E.coli,S115-S270

Cross-Reactivity (Details): There is no detectable cross-reactivity with other relevant proteins.

Sensitivity: <0.5pg/mL

Product Details

Material not included: Microplate reader in standard size. Automated plate washer. Adjustable pipettes and pipette tips. Multichannel pipettes are recommended in the condition of large amount of samples in the detection. Clean tubes and Eppendorf tubes. Washing buffer (neutral PBS or TBS). Preparation of 0.01M TBS: Add 1.2g Tris, 8.5g NaCl

Target Details

Target: IL1A

Alternative Name: IL1A ([IL1A Products](#))

Background: Protein Function: Produced by activated macrophages, IL-1 stimulates thymocyte proliferation by inducing IL-2 release, B-cell maturation and proliferation, and fibroblast growth factor activity. IL-1 proteins are involved in the inflammatory response, being identified as endogenous pyrogens, and are reported to stimulate the release of prostaglandin and collagenase from synovial cells.

Background: Interleukin-1 alpha(IL-1 alpha) and interleukin-1 beta(IL-1 beta) are two biochemically distinct, but distantly related, polypeptidic cytokines that play a key role in inflammation, immunologic reactions, and tissue repair. IL-1 alpha has been implicated in the pathogenesis of infectious, autoimmune and inflammatory diseases. Recently, it has been shown that IL-1 alpha is identical to hematopoietin 1, which is described as a hematopoietic growth factor acting on early progenitor cells in synergy with other hematopoietic growth factors.¹ The human interleukin 1 alpha gene is assigned to chromosome 2. Genetic polymorphisms at interleukin(IL)-1alpha and IL-1beta have been recently suggested to be associated with severity of adult periodontitis. The murine IL-1 alpha and IL-1 beta genes encode structurally and evolutionarily related cytokines that exert a regulatory role in numerous physiological processes including hemopoiesis. The standard product used in this kit is recombinant rat IL-1alpha with the molecular mass of 18 kDa.

Synonyms: Interleukin-1 alpha,IL-1 alpha,IL1a,

Full Gene Name: Interleukin-1 alpha

Cellular Localisation: Secreted. The lack of a specific hydrophobic segment in the precursor sequence suggests that IL-1 is released by damaged cells or is secreted by a mechanism differing from that used for other secretory proteins.

Gene ID: 24493

UniProt: [P16598](#)

Pathways: [NF-kappaB Signaling](#), [Autophagy](#), [Cancer Immune Checkpoints](#)

Application Details

Application Notes:	Before using Kit, spin tubes and bring down all components to bottom of tube. Duplicate well assay was recommended for both standard and sample testing.
Plate:	Pre-coated
Protocol:	rat IL-1 alpha ELISA Kit was based on standard sandwich enzyme-linked immune-sorbent assay technology. A monoclonal antibody from mouse specific for IL-1 alpha has been precoated onto 96-well plates. Standards (E.coli,S115-S270) and test samples are added to the wells, a biotinylated detection polyclonal antibody from goat specific for IL-1 alpha is added subsequently and then followed by washing with PBS or TBS buffer. Avidin-Biotin-Peroxidase Complex was added and unbound conjugates were washed away with PBS or TBS buffer. HRP substrate TMB was used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the rat IL-1 alpha amount of sample captured in plate.
Assay Procedure:	Aliquot 0.1 mL per well of the 300pg/mL, 150pg/mL, 75pg/mL, 37.5pg/mL, 18.8pg/mL, 9.4pg/mL, 4.69pg/mL rat IL-1 alpha standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of rat cell culture supernates or serum to each empty well. See "Sample Dilution Guideline" above for details. We recommend that each rat IL-1 alpha standard solution and each sample is measured in duplicate.
Assay Precision:	<ul style="list-style-type: none">• Sample 1: n=16, Mean(pg/ml): 45, Standard deviation: 1.89, CV(%): 4.2• Sample 2: n=16, Mean(pg/ml): 137, Standard deviation: 6.58, CV(%): 4.8• Sample 3: n=16, Mean(pg/ml): 211, Standard deviation: 10.76, CV(%): 5.1,• Sample 1: n=24, Mean(pg/ml): 49, Standard deviation: 2.646, CV(%): 5.4• Sample 2: n=24, Mean(pg/ml): 145, Standard deviation: 8.99, CV(%): 6.2• Sample 3: n=24, Mean(pg/ml): 230, Standard deviation: 15.41, CV(%): 6.7
Restrictions:	For Research Use only

Handling

Handling Advice:	Avoid multiple freeze-thaw cycles.
Storage:	-20 °C,4 °C
Storage Comment:	Store at 4°C for 6 months, at -20°C for 12 months. Avoid multiple freeze-thaw cycles
Expiry Date:	12 months

Publications

Product cited in:


Sai, Yao, Shen, Zheng, Sun, Wu, Wang, Yao: "Dynamic expression of hepatic GP73 mRNA and protein and circulating GP73 during hepatocytes malignant transformation." in: **Hepatobiliary & pancreatic diseases international : HBPD INT**, Vol. 19, Issue 5, pp. 449-454, (2020) ([PubMed](#)).

Dong, Chen, Li, Li, Wen, Lin, Ma, Wei, Chen, Ruan, Lin, Wang, Wu, Wu: "Serum Golgi protein 73 is a prognostic rather than diagnostic marker in hepatocellular carcinoma." in: **Oncology letters**, Vol. 14, Issue 5, pp. 6277-6284, (2017) ([PubMed](#)).

Kosanam, Prassas, Chrystoja, Soleas, Chan, Dimitromanolakis, Blasutig, Rückert, Gruetzmann, Pilarsky, Maekawa, Brand, Diamandis: "Laminin, gamma 2 (LAMC2): a promising new putative pancreatic cancer biomarker identified by proteomic analysis of pancreatic adenocarcinoma tissues." in: **Molecular & cellular proteomics : MCP**, Vol. 12, Issue 10, pp. 2820-32, (2013) ([PubMed](#)).

There are more publications referencing this product on: [Product page](#)

Images



ELISA

Image 1. Rat IL-1 alpha PicoKine ELISA Kit standard curve