

Datasheet for ABIN411332

**MMP8 ELISA Kit**[Go to Product page](#)**1** Image**3** Publications

## Overview

|                          |                 |
|--------------------------|-----------------|
| Quantity:                | 96 tests        |
| Target:                  | MMP8            |
| Binding Specificity:     | AA 21-467       |
| Reactivity:              | Human           |
| Method Type:             | Sandwich ELISA  |
| Detection Range:         | 156-10000 pg/mL |
| Minimum Detection Limit: | 156 pg/mL       |
| Application:             | ELISA           |

## Product Details

|                             |   |
|-----------------------------|---|
| Purpose:                    | Sandwich High Sensitivity ELISA kit for Quantitative Detection of Human MMP-8 |
| Brand:                      | PicoKine™   |
| Sample Type:                | Cell Culture Supernatant, Serum, Plasma (heparin), Saliva                     |
| Analytical Method:          | Quantitative  |
| Detection Method:           | Colorimetric  |
| Immunogen:                  | Expression system for standard: NSO<br>Immunogen sequence: F21-G467           |
| Specificity:                | Expression system for standard: NSO<br>Immunogen sequence: F21-G467           |
| Cross-Reactivity (Details): | There is no detectable cross-reactivity with other relevant proteins.         |

## Product Details

|                        |  |
|------------------------|--|
| Sensitivity:           | <10pg/mL   |
| Material not included: | Microplate reader in standard size. Automated plate washer. Adjustable pipettes and pipette tips. Multichannel pipettes are recommended in the condition of large amount of samples in the detection. Clean tubes and Eppendorf tubes. Washing buffer (neutral PBS or TBS). Preparation of 0.01M TBS: Add 1.2g Tris, 8.5g NaCl |

## Target Details

|                   |  |
|-------------------|--|
| Target:           | MMP8   |
| Alternative Name: | MMP8 ( <a href="#">MMP8 Products</a> )   |
| Background:       | <p>Protein Function: Can degrade fibrillar type I, II, and III collagens.</p> <p>Background: Matrix metalloproteinase 8(MMP8) also called neutrophil collagenase. Neutrophil collagenase, a member of the family of matrix metalloproteinases, is distinct from the collagenase of skin fibroblasts and synovial cells in substrate specificity and immunologic crossreactivity. MMP8, an enzyme that degrades fibrillar collagens imparting strength to the fetal membranes, is expressed by leukocytes and chorionic cytotrophoblast cells. The human neutrophil collagenase(HNC) cDNA clone has been sequenced and shown to encode a 467-residue protein. Neutrophil collagenase has been found to possess 57 % identity with the deduced protein sequence for fibroblast collagenase with 72 % chemical similarity. Certain regions of the molecule, including the putative zinc-binding region, are highly conserved. When compared with the published sequence for fibroblast collagenase, neutrophil collagenase contains four additional sites for glycosylation. The standard product used in this kit is natural, isolating from human MMP-8. The detected MMP-8 includes zymogen and active enzyme.</p> <p>Synonyms: Neutrophil collagenase,3.4.24.34,Matrix metalloproteinase-8,MMP-8,PMNL collagenase,PMNL-CL,MMP8,CLG1,</p> <p>Full Gene Name: Neutrophil collagenase</p> <p>Cellular Localisation: Cytoplasmic granule. Secreted, extracellular space, extracellular matrix . Stored in intracellular granules.</p> |
| Gene ID:          | 4317   |
| UniProt:          | <a href="#">P22894</a>   |

## Application Details

|                    |  |
|--------------------|--|
| Application Notes: | Before using Kit, spin tubes and bring down all components to bottom of tube. Duplicate well assay was recommended for both standard and sample testing. |
|--------------------|--|

## Application Details

|                  |   |
|------------------|---|
| Comment:         | Sequence similarities: Belongs to the peptidase M10A family.<br>Tissue Specificity: Neutrophils.  |
| Plate:           | Pre-coated  |
| Protocol:        | human MMP-8 ELISA Kit was based on standard sandwich enzyme-linked immune-sorbent assay technology. A monoclonal antibody from mouse specific for MMP-8 has been precoated onto 96-well plates. Standards(NSO, F21-G467) and test samples are added to the wells, a biotinylated detection polyclonal antibody from goat specific for MMP-8 is added subsequently and then followed by washing with PBS or TBS buffer. Avidin-Biotin-Peroxidase Complex was added and unbound conjugates were washed away with PBS or TBS buffer. HRP substrate TMB was used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the human MMP-8 amount of sample captured in plate. |
| Assay Procedure: | Aliquot 0.1 mL per well of the 10000pg/mL, 5000pg/mL, 2500pg/mL, 1250pg/mL, 625pg/mL, 312pg/mL, 156pg/mL human MMP-8 standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of human cell culture supernates, serum, plasma (heparin) or saliva to each empty well. See "Sample Dilution Guideline" above for details. It is recommended that each human MMP-8 standard solution and each sample be measured in duplicate.  |
| Assay Precision: | <ul style="list-style-type: none"><li>• Sample 1: n=16, Mean(pg/ml): 1275, Standard deviation: 56.1, CV(%): 4.4</li><li>• Sample 2: n=16, Mean(pg/ml): 3563, Standard deviation: 185.3, CV(%): 5.2</li><li>• Sample 3: n=16, Mean(pg/ml): 6822, Standard deviation: 395.7, CV(%): 5.8,</li><li>• Sample 1: n=24, Mean(pg/ml): 1370, Standard deviation: 69.87, CV(%): 5.1</li><li>• Sample 2: n=24, Mean(pg/ml): 3728, Standard deviation: 242.32, CV(%): 6.5</li><li>• Sample 3: n=24, Mean(pg/ml): 7142, Standard deviation: 521.4, CV(%): 7.3</li></ul>  |
| Restrictions:    | For Research Use only   |

## Handling

|                  |  |
|------------------|--|
| Handling Advice: | Avoid multiple freeze-thaw cycles.   |
| Storage:         | -20 °C, 4 °C   |
| Storage Comment: | Store at 4°C for 6 months, at -20°C for 12 months. Avoid multiple freeze-thaw cycles |
| Expiry Date:     | 12 months  |

Publications

Product cited in: Wang, Lee, Chou, Yang, Wei, Chen, Yao, Hsu, Zhu, Ying, Ye, Wang, Lim, Xia, Ko, Liu, Liu, Wu, Wang, Li, Prakash, Katz, Kang, Kim, Fleming, Fogelman, Javle, Maitra, Hung: "Angiogenin/Ribonuclease 5 Is an EGFR Ligand and a Serum Biomarker for Erlotinib Sensitivity in Pancreatic Cancer." in: **Cancer cell**, Vol. 33, Issue 4, pp. 752-769.e8, (2019) ([PubMed](#)).

Images

