

Datasheet for ABIN4368347

Recombinant anti-Blue Fluorescent Protein antibody (Atto 488)



Go to Product page

| (| ۱۱/ | e | r\/ | Ì١ | | ۱۸ | |
|---|-----|--------|-----|----|---------------|----|---|
| | , v | \cup | V | 1 | $\overline{}$ | V | V |

Cross-Reactivity (Details):

Characteristics:

| Quantity: | 200 μL | |
|--------------------|---|--|
| Target: | Blue Fluorescent Protein (BFP) | |
| Reactivity: | Entacmaea quadricolor | |
| Host: | Alpaca | |
| Expression System: | E.coli | |
| Antibody Type: | Recombinant Antibody | |
| Clonality: | Monoclonal | |
| Conjugate: | This Blue Fluorescent Protein antibody is conjugated to Atto 488 | |
| Application: | Flow Cytometry (FACS), Immunocytochemistry (ICC), Immunofluorescence (IF) | |
| Product Details | | |
| Purpose: | Camelid sdAb anti-TagFP conjugated with Atto488, Clone 1H7 | |
| Immunogen: | BFP | |
| Clone: | 1H7 | |
| Fragment: | single-domain Antibody (sdAb) | |
| Specificity: | Recognizes mTagBFP, mKate, mKate2, mTagRFP, mTagRFP657 and most common fluorescent proteins deriving from Entacmaea quadricolor | |
| | | |

Does not cross-react with common GFP- or dsRed derivatives.

A dye conjugated alpaca single-domain antibody (sdAb), also referred to as VHH or nanobody.

It is in the range of 15 kDa and 3 nm in molecular weight and size respectively. This means that

Product Details

| 1 Toddot Betailo | |
|---------------------|--|
| | sdAbs are ten times lighter and up to 5x smaller than a conventional IgG molecule. They can |
| | position a fluorophore up to 20 nm closer to the intended target than using conventional primary-secondary antibody complex detection. |
| Purification: | Produced in: E.coli |
| Labeling Ratio: | a single fluorophore is coupled to exactly one sdAb |
| T | |
| Target Details | |
| Target: | Blue Fluorescent Protein (BFP) |
| Alternative Name: | TagBFP (BFP Products) |
| Molecular Weight: | 26 kDa |
| Application Details | |
| Application Notes: | Recommended dilution 1:500 |
| Comment: | Each fluorophore is coupled to exactly one sdAb, which in turn binds to its target molecule in a monovalent fashion. The high binding affinity and a high coupling efficiency of > 95% |
| | guarantees a highly linear relation between target molecule number and fluorescent intensity. |
| | This enables you to directly count your target molecule of interest. The fluorophore is located |
| | exceptionally close to the recognized epitope (< 1.5 nm), which is ideal for all microscopy |
| | techniques. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Lyophilized |
| Reconstitution: | Reconstitute with 50 % glycerol in deionized water. We recommend including 0.1 % sodium |
| | azide as a preservative if applicable. |
| Concentration: | 5 μΜ |
| Buffer: | lyophilized from PBS pH7.4 with 2% BSA (US-Origin) |
| Handling Advice: | Protect from light! |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Up to 3 months store at -20 °C. Up to 12 months store at -80 °C or below |
| | |

| | 1 | l: |
|---|------|--------|
| - | ココロロ | urnen |
| | Hand | III IU |
| | | |

Expiry Date:

12 months