antibodies.com

## Datasheet for ABIN4371359 ULBP1 Protein (AA 26-211) (His tag)



| Quantity:                     | 50 µg   |
|-------------------------------|---|
| Target:                       | ULBP1   |
| Protein Characteristics:      | AA 26-211   |
| Origin:                       | Mouse   |
| Source:                       | Human Cells   |
| Protein Type:                 | Recombinant   |
| Purification tag / Conjugate: | This ULBP1 protein is labelled with His tag.                      |
| Product Details               |   |
| Purpose:                      | Recombinant Mouse NKG2D Ligand 1/NKG2DL/ULBP1 (C-6His)            |
| Sequence:                     | PRIEETASLC NIYKVNRSES GQHSHEVQGL LNRQPLFVYK DKKCHAIGAH RNSMNATKIC |
|                               | EKEVDTLKDG IDIFKGLLLH IVQETNTTGK PLTLQAEVCG QYEVDKHFTG YAIVSLNGKN |
|                               | IFRVDTSTGN WTQLDHEFEK FIEMCKEDKV LAAFLKKTTE GDCRTWLDEL MLHWKEHLEF |
|                               | AGSFSTVDHH HHHH   |
| Characteristics:              | Recombinant Mouse NKG2D Ligand 1/NKG2DL/ULBP1 (C-6His)            |
| Purity:                       | > 95 % as determined by reducing SDS-PAGE.                        |
| Sterility:                    | 0.2 µm filtered   |
| Endotoxin Level:              | Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test          |
| Target Details                |   |
| Target:                       | ULBP1   |

Page 1/2 | Product datasheet for ABIN4371359 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

| Target Details   |   |
|--|---|
| Alternative Name:  | NKG2D Ligand 1 (ULBP1 Products)   |
| Molecular Weight:  | 22.3 kDa  |
| UniProt:   | Q8HWA3  |
| Pathways:  | Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process   |
| Application Details  |   |
| Restrictions:  | For Research Use only   |
| Handling   |   |
|  |   |
| Format:  | Lyophilized   |
| Format:<br>Reconstitution:                                 | Lyophilized<br>It is not recommended to reconstitute to a concentration less than 100 µg/mL.  |
|  |   |
|  | It is not recommended to reconstitute to a concentration less than 100 µg/mL.   |
|  | It is not recommended to reconstitute to a concentration less than 100 µg/mL.<br>Dissolve the lyophilized protein in ddH20.   |
| Reconstitution:  | It is not recommended to reconstitute to a concentration less than 100 µg/mL.<br>Dissolve the lyophilized protein in ddH20.<br>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.  |
| Reconstitution:<br>Buffer:                                 | It is not recommended to reconstitute to a concentration less than 100 µg/mL.<br>Dissolve the lyophilized protein in ddH2O.<br>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.<br>Lyophilized from a 0.2 µm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.   |
| Reconstitution:<br>Buffer:<br>Handling Advice:             | It is not recommended to reconstitute to a concentration less than 100 µg/mL.<br>Dissolve the lyophilized protein in ddH2O.<br>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.<br>Lyophilized from a 0.2 µm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.<br>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.                       |
| Reconstitution:<br>Buffer:<br>Handling Advice:<br>Storage: | It is not recommended to reconstitute to a concentration less than 100 µg/mL.<br>Dissolve the lyophilized protein in ddH2O.<br>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.<br>Lyophilized from a 0.2 µm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.<br>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.<br>4 °C/-20 °C/-80 °C |