



[Go to Product page](#)

Datasheet for ABIN457262

Prothrombin Fragment 1+2 ELISA Kit

1 Image

1 Publication

Overview

| | |
|--------------------------|---------------------------------|
| Quantity: | 96 tests |
| Target: | Prothrombin Fragment 1+2 (F1+2) |
| Binding Specificity: | Fragment 1, Fragment 2 |
| Reactivity: | Rat |
| Method Type: | Sandwich ELISA |
| Detection Range: | 62.5-4000 pM/L |
| Minimum Detection Limit: | 62.5 pM/L |
| Application: | ELISA |

Product Details

| | |
|-----------------------------|--|
| Purpose: | For the quantitative determination of rat prothrombin fragment 1+2 (F1+2) concentrations in serum, plasma and tissue homogenates. |
| Sample Type: | Serum, Plasma, Tissue Homogenate |
| Analytical Method: | Quantitative |
| Detection Method: | Colorimetric |
| Specificity: | This assay has high sensitivity and excellent specificity for detection of rat F1+2. |
| Cross-Reactivity (Details): | Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between the target antigen and all analogues for other species. Therefore, cross reaction may still exist. |
| Sensitivity: | 15.6 pM/L |

Product Details

- Components:
- Assay plate (12 × 8 coated Microwells)
 - Standard (freeze dried)
 - Biotin-antibody (100 × concentrate)
 - HRP-avidin (100 × concentrate)
 - Biotin-antibody Diluent
 - HRP-avidin Diluent
 - Sample Diluent
 - Wash Buffer (25 × concentrate)
 - TMB Substrate
 - Stop Solution
 - Adhesive Strip (for 96 wells)
 - Instruction manual

Target Details

- Target: Prothrombin Fragment 1+2 (F1+2)
- Alternative Name: prothrombin fragment 1+2,F1+2 ([F1+2 Products](#))

Application Details

- Application Notes:
- The supplier is only responsible for the kit itself, but not for the samples consumed during the assay. The user should calculate the possible amount of the samples used in the whole test. Please reserve sufficient samples in advance.
 - Samples to be used within 5 days may be stored at 2-8°C, otherwise samples must be stored at -20°C (≤ 1 month) or -80°C (≤ 2 months) to avoid loss of bioactivity and contamination.
 - Grossly hemolyzed samples are not suitable for use in this assay.
 - If the samples are not indicated in the manual, a preliminary experiment to determine the validity of the kit is necessary.
 - Please predict the concentration before assaying. If values for these are not within the range of the standard curve, users must determine the optimal sample dilutions for their particular experiments.
 - Tissue or cell extraction samples prepared by chemical lysis buffer may cause unexpected ELISA results due to the impacts of certain chemicals.
 - Owing to the possibility of mismatching between antigens from another resource and antibodies used in this supplier's kits (e.g., antibody targets conformational epitope rather than linear epitope), some native or recombinant proteins from other manufacturers may not be recognized by this supplier's products.
 - Influenced by factors including cell viability, cell number and cell sampling time, samples from cell culture supernatant may not be recognized by the kit.
 - Fresh samples without long time storage are recommended for the test. Otherwise, protein degradation and denaturalization may occur in those samples and finally lead to wrong results.

Application Details

Comment: Detection wavelength: 450 nm

Information on standard material:

Depending on the antigen to be detected, standards can be either native or recombinant protein. The recombinant proteins are being expressed in CHO cells in most cases. Please inquire for more information. The formulation of auxiliary material in the standard is considered proprietary information, however it does not contain any poisonous substance. Proclin 300 (1:3000) is used as preservative.

Information on reagents:

In most cases the stop solution provided is 1 N H₂SO₄. The formulation of wash solution is proprietary information. None of the components contain (sodium) azide, thimerosal, 2-mercaptoethanol (2-ME) or any other poisonous materials. For the sandwich method kits, the sample diluent, antibody diluent, enzyme diluent and standard all contain BSA.

Information on antibodies:

The antibodies provided in different kits vary in regards to clonality and host. Some antibodies are affinity purified, some are Protein A

Sample Volume: 100 µL

Assay Time: 1 - 4.5 h

Plate: Pre-coated

Protocol: This assay employs the quantitative sandwich enzyme immunoassay technique. Antibody specific for F1+2 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any F1+2 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for F1+2 is added to the wells. After washing, avidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of F1+2 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Assay Precision: Intra-assay precision (precision within an assay): Three samples of known concentration were tested twenty times on one plate to assess precision.

Inter-assay precision (precision between assays): Three samples of known concentration were tested in twenty assays to assess precision.

Application Details

- Intra-assay: CV% less than 8%
- Inter-assay: CV% less than 10%

Restrictions: For Research Use only

Handling

Precaution of Use: The Stop Solution provided with this kit is an acid solution. Wear eye, hand, face and clothing protection when using this material.

Handling Advice:

- The kit should not be used beyond the expiration date on the kit label.
- Do not mix or substitute reagents with those from other lots or sources.
- If samples generate values higher than the highest standard, dilute the samples with Sample Diluent and repeat the assay.
- Any variation in Sample Diluent, operator, pipetting technique, washing technique, incubation time/temperature and kit age can cause variation in binding.
- This assay is designed to eliminate interference by soluble receptors, binding proteins and other factors present in biological samples. Until all factors have been tested in the Immunoassay, the possibility of interference cannot be excluded.

Storage: 4 °C/-20 °C

Storage Comment: For unopened kit: All the reagents should be kept according to the labels on vials.

Expiry Date: 6 months


Publications

Product cited in: Mboumba Bouassa, Péré, Gubavu, Prazuck, Jenabian, Veyer, Meye, Touzé, Bélec: "Serum and cervicovaginal IgG immune responses against $\alpha 7$ and $\alpha 9$ HPV in non-vaccinated women at risk for cervical cancer: Implication for catch-up prophylactic HPV vaccination." in: **PLoS ONE**, Vol. 15, Issue 5, pp. e0233084, (2020) ([PubMed](#)).

Bystrická, Zatovicová, Petříková, Soláriková, Russ, Ziegler: "Monoclonal antibodies suitable for type-specific identification of herpes simplex viruses by a rapid culture assay." in: **Acta virologica**, Vol. 43, Issue 6, pp. 399-402, (2000) ([PubMed](#)).

Bystrická, Petříková, Zatovicová, Soláriková, Kostolanský, Mucha, Russ: "Monoclonal antibodies to the distinct antigenic sites on glycoproteins C and B and their protective abilities in herpes simplex virus infection." in: **Acta virologica**, Vol. 41, Issue 1, pp. 5-12, (1997) ([PubMed](#)).

Bystrická, Vancíková, Kasalová, Rajcáni, Kostál, Murányiová, Poláková, Russ: "Type-common and type-specific monoclonal antibodies to herpes simplex virus types 1 and 2." in: **Acta virologica**, Vol. 35, Issue 2, pp. 152-64, (1991) ([PubMed](#)).



ELISA

Image 1. Typical standard curve