


Datasheet for ABIN457304

anti-E-cadherin antibody (APC)





Go to Product page

_					
()	VE	۲۱د	/1/	\square	٨.

Quantity:	100 tests	
Target:	E-cadherin (CDH1)	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This E-cadherin antibody is conjugated to APC	
Application:	Flow Cytometry (FACS)	

Product Details

Purpose:	Anti-Hu CD324 APC	
Immunogen:	T-47D cells	
Clone:	67A4	
Isotype:	lgG1	
Specificity:	The mouse monoclonal antibody 67A4 recognizes an extracellular epitope of CD324 / E-cadherin, an approximately 100 kDa epithelial cell adhesion molecule, whose detection is important for determination of invasive potential of epithelial neoplasms.	
Cross-Reactivity (Details):	ails): Human	
Purification:	Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.	

Target Details

Target:	E-cadherin (CDH1)	
Alternative Name:	CD324 (CDH1 Products)	
Background:	Cadherin 1,CD324 / E-cadherin is an epithelial cell surface molecule, which provides calcium-dependent homophilic interactions with E-cadherin of another cell. These intaractions take part in morphogenetic programs controlling the maintenance of the structural and functional integrity of epithelia and affect invasive potential of epithelial neoplasms. CD324 / E-cadherin is implicated in cell growth and differentiation, cell recognition, and sorting during developmental morphogenesis, as well as in aggregation-dependent cell survival. CD324 / E-cadherin-mediated cell adhesion system is highly regulated from inside the cell by a number of intracellular signaling pathways.,Cadherin-1, E-cadherin, Uvomorulin, CAM 120/80, UVO, CDHE, ECAD, LCAM, BCDS1, Arc-1	
Gene ID:	999	
UniProt:	P12830	
Pathways:	WNT Signaling, Sensory Perception of Sound, Cell-Cell Junction Organization, Tube Formation	
Application Details		
Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 10 μ L reagent / 100 μ L of whole blood or 10 ⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.	
Restrictions:	For Research Use only	
Handling		
Reconstitution:	No reconstitution is necessary.	
Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Handling Advice:	Do not freeze. Avoid prolonged exposure to light.	
Storage:	4 °C	
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.	

Product cited in:

Lin, Liao, Lee, Hung, Sayion, Chen, Kang, Huang, Cherng: "Molecular events associated with epithelial to mesenchymal transition of nasopharyngeal carcinoma cells in the absence of Epstein-Barr virus genome." in: **Journal of biomedical science**, Vol. 16, pp. 105, (2009) (PubMed).

Caberg, Hubert, Begon, Herfs, Roncarati, Boniver, Delvenne: "Silencing of E7 oncogene restores functional E-cadherin expression in human papillomavirus 16-transformed keratinocytes." in: **Carcinogenesis**, Vol. 29, Issue 7, pp. 1441-7, (2008) (PubMed).


Furio, Guezennec, Ducarre, Guesnet, Peguet-Navarro: "Differential effects of allergens and irritants on early differentiating monocyte-derived dendritic cells." in: **European journal of dermatology: EJD**, Vol. 18, Issue 2, pp. 141-7, (2008) (PubMed).

Robertson, Ali, McDonnell, Burt, Kirby: "Chronic renal allograft dysfunction: the role of T cell-mediated tubular epithelial to mesenchymal cell transition." in: **Journal of the American Society of Nephrology: JASN**, Vol. 15, Issue 2, pp. 390-7, (2004) (PubMed).

Kutlesa, Wessels, Speiser, Steiert, Müller, Klein: "E-cadherin-mediated interactions of thymic epithelial cells with CD103+ thymocytes lead to enhanced thymocyte cell proliferation." in: **Journal of cell science**, Vol. 115, Issue Pt 23, pp. 4505-15, (2002) (PubMed).

There are more publications referencing this product on: Product page

Images



Flow Cytometry

Image 1. Separation of HT-29 cell line (red) from SP2 cell line (blue) in flow cytometry analysis (surface staining) stained using anti-human CD324 (67A4) APC antibody.