.-online.com antibodies

Datasheet for ABIN457421 anti-CD86 antibody

2 Images

7 Publications



Overview

Quantity:	0.1 mg
Target:	CD86
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Application:	Flow Cytometry (FACS), Immunoprecipitation (IP), Immunohistochemistry (Frozen Sections)
	(IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	LPS-activated CBA/Cs mouse splenic B cells
Clone:	GL-1
lsotype:	IgG2a kappa
Specificity:	The rat monoclonal antibody GL-1 reacts with an extracellular epitope of CD86 (B7-2), a 70-80 kDa type I transmembrane glycoprotein of immunoglobulin supergene family, expressed on professional antigen-presenting cells, such as dendritic cells, macrophages or activated B lymphocytes.
Cross-Reactivity (Details):	Mouse
Purification:	Purified by protein-G affinity chromatography.
Purity:	> 95 % (by SDS-PAGE)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN457421 | 02/07/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Target:	CD86
Alternative Name:	CD86 (CD86 Products)
Background:	CD86 Molecule,CD80 (B7-1) and CD86 (B7-2) are ligands of T cell critical costimulatory molecule CD28 and of an inhibitory receptor CTLA-4 (CD152). The both B7 Molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, the both molecules can also substitute for each other in this process. The question what are the differences in CD80 and CD86 competency has not been fully elucidated yet, there are still conflicts in results about their respective roles in initiation or sustaining of the T cell immune response.,B7-2, FUN-1, Ly58
Gene ID:	12524
UniProt:	P42082
Pathways:	TCR Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Activation of Innate immune Response, Cellular Response to Molecule of Bacterial Origin, Positive Regulation of Immune Effector Process, Activated T Cell Proliferation

Application Details

Application Notes:	Flow cytometry: Recommended dilution: 2 µg/mL, positive control: murine splenocytes.
Restrictions:	For Research Use only
Handling	
Concentration:	1 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Do not freeze.

Product cited in:

Nolan, Kobayashi, Naveed, Kelly, Hoshino, Hoshino, Karulf, Rom, Weiden, Gold: "Differential role for CD80 and CD86 in the regulation of the innate immune response in murine polymicrobial sepsis." in: **PLoS ONE**, Vol. 4, Issue 8, pp. e6600, (2009) (PubMed).


Edgtton, Kausman, Li, OSullivan, Lo, Hutchinson, Yagita, Holdsworth, Kitching: "Intrarenal antigens activate CD4+ cells via co-stimulatory signals from dendritic cells." in: **Journal of the American Society of Nephrology : JASN**, Vol. 19, Issue 3, pp. 515-26, (2008) (PubMed).

Nolan, Weiden, Kelly, Hoshino, Hoshino, Mehta, Gold: "CD40 and CD80/86 act synergistically to regulate inflammation and mortality in polymicrobial sepsis." in: **American journal of respiratory and critical care medicine**, Vol. 177, Issue 3, pp. 301-8, (2008) (PubMed).

Steptoe, Ritchie, Jones, Harrison: "Autoimmune diabetes is suppressed by transfer of proinsulin-encoding Gr-1+ myeloid progenitor cells that differentiate in vivo into resting dendritic cells." in: **Diabetes**, Vol. 54, Issue 2, pp. 434-42, (2005) (PubMed).

Brasel, De Smedt, Smith, Maliszewski: "Generation of murine dendritic cells from flt3-ligandsupplemented bone marrow cultures." in: **Blood**, Vol. 96, Issue 9, pp. 3029-39, (2000) (PubMed).

There are more publications referencing this product on: Product page




Flow Cytometry

Image 1. Flow cytometry surface staining pattern of murine peritoneal fluid cells suspension stained using anti-mouse CD86 (GL-1) purified antibody (concentration in sample 0,6 µg/mL) DAR APC.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN457421 | 02/07/2024 | Copyright antibodies-online. All rights reserved.

Images

Images



Flow Cytometry

Image 2. Separation of murine CD86 positive myeloid cells (red-filled) from murine CD86 negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of murine peritoneal fluid cells suspension stained using antimouse CD86 (GL-1) purified antibody (concentration in sample 0,6 μ g/mL) DAR APC.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/4 | Product datasheet for ABIN457421 | 02/07/2024 | Copyright antibodies-online. All rights reserved.