

Datasheet for ABIN457556

anti-HCV 1b NS5B antibody (AA 77-86)**2** Images[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	HCV 1b NS5B (HCV NS5b)
Binding Specificity:	AA 77-86
Reactivity:	Hepatitis C Virus (HCV)
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This HCV 1b NS5B antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunoprecipitation (IP)

Product Details

Immunogen:	RecombinantHepatitis C Virus subtype 1b nonstructural protein 5B (NS5B) RNA-dependent RNApolymerase (RdRp)
Sequence:	TVKAKLLSVE
Clone:	6G5
Isotype:	IgG1
Specificity:	Human, HCV subtype 1b NS5B
Purification:	purified by protein G affinity chromatography

Target Details

Target:	HCV 1b NS5B (HCV NS5b)
---------	------------------------

Target Details


Abstract:	HCV NS5b Products
Target Type:	Viral Protein
Background:	Non-structural protein 5B (NS5B) represents the RNA-dependent RNA polymerase (RdRp) of Hepatitis C Virus, which is a small positive strand RNA virus in the family Flaviviridae. HCV is a major causative agent of acute and chronic hepatitis, hepatocellular carcinoma and liver cirrhosis. The single subunit RNA-dependent RNA polymerase is absolutely essential for the viral replication.
UniProt:	Q9WMX2

Application Details

Application Notes:	ELISA titer 1:1000, Western immunoblotting, reduced conditions, dilution 1:1000, Immunoprecipitation. Monoclonal antibody working titer has to be established practically for each particular antigen and assay format.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/ml
Buffer:	PBS pH 7.4, with 0.1%
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Divide antibody into aliquots prior usage. Avoid multiple freeze-thaw cycles as product degradation may result.
Storage:	-20 °C
Storage Comment:	This product is shipped in non-frozen liquid form in ambient conditions



Immunoprecipitation

Image 1. IP was carried out with NS5B specific mAb 6G5 using the lysates of Huh7 cells harboring selectable subgenomic HCV RNA replicon (upper panel) or plain Huh7 cells (lower panel). NS5B polyclonal antibodies (α-NS5B) and α-Bcl2 mAb, directed against cellular protein, were used as positive and negative controls respectively. Asterisk indicates immunoglobulin heavy chain; protein blots were probed with NS5B specific mAb 10D6.

Western Blotting

Image 2. WB analysis of Huh-7 cell line expressing NS5B protein. Line 1. Huh-7 transfected with HCV replicon, Line 2. Huh-7 non-transfected cells.