

Datasheet for ABIN457687 **anti-Albumin antibody**



[Go to Product page](#)

Overview

Quantity:	1 mL
Target:	Albumin (ALB)
Reactivity:	Duck
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This Albumin antibody is un-conjugated
Application:	ELISA, Immunoelectrophoresis (IEP), Immunodiffusion (ID), Radial Immunodiffusion (RID), Nephelometry (Neph)

Product Details

Immunogen:	Highly purified albumin isolated from duck serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Specificity:	Inter-species cross-reactivity is a normal feature of antibodies to serum proteins, since homologous proteins of different species frequently share antigenic determinants. The degree of cross-reactivity is also dependent on the concentrations of the reactants and the sensitivity of the assay arrangement. This antiserum has not been tested for cross-reactivity.
Characteristics:	Precipitating polyclonal goat antiserum to duck serum albumin
Purification:	Adsorption: Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies reacting with other serum proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.

Target Details

Target:	Albumin (ALB)
Alternative Name:	Serum Albumin (ALB Products)
Background:	The defined antibody specificity is directed to albumin as tested against duck sera. In immunoelectro- phoresis and double radial immunodiffusion (Ouchterlony), using various antiserum concentrations against appropriate concentrations of the immunogen, a single characteristic precipitin line is obtained which shows a reaction of identity with the precipitin lines obtained against duck serum and the purified albumin.
Pathways:	Lipid Metabolism

Application Details

Application Notes:	As precipitating antiserum to identify or measure duck albumin by a variety of immunodiffusion techniques, including immunoelectrophoresis, single and double radial immunodiffusion (Mancini, Ouchterlony) and electroimmunodiffusion (Laurell). It has not been tested for use in nephelometry, ELISA or immunochemistry, but this does not exclude such use if proper controls are included.
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Concentration:	Total protein and IgG concentrations in the antiserum are comparable to those in pooled goat serum. No foreign proteins added. Antibody titre: Precipitin titre not less than 1:64 when tested against normal duck serum in agar block immunodiffusion titratio
Buffer:	Delipidated, heat inactivated, lyophilized, stable whole antiserum
Preservative:	Without preservative
Storage:	4 °C/-20 °C
Storage Comment:	The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C, prolonged storage at or below -20°C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day.