



[Go to Product page](#)

Datasheet for ABIN457747

Goat anti-Guinea Pig IgG (Fc Region) Antibody

Overview

Quantity:	1 mL
Target:	IgG
Binding Specificity:	Fc Region
Reactivity:	Guinea Pig
Host:	Goat
Clonality:	Polyclonal
Application:	Immunoprecipitation (IP)

Product Details

Immunogen:	Highly purified normal IgG isolated from pooled guinea pig serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Specificity:	Precipitating polyclonal Goat antiSerum to Guinea Pig IgG, Fc specific
Cross-Reactivity (Details):	The antiSerum does not cross-react with any other component of the Guinea Pig immunoglobulin system. Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Crossreactivity of this antiSerum has not been tested in detail.
Characteristics:	The reactivity of the antiserum is restricted to the Fc part of both subclasses IgG1 and IgG2. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal guinea pig serum a single precipitin line is obtained which shows a reaction of identity with the precipitin line obtained with purified IgG. No precipitation reaction is obtained with purified IgA, IgM, and IgG/Fab fragments. In precipitating techniques as immunoelectrophoresis and radial immunodiffusion to identify the presence of IgG in guinea

Product Details

pig serum or other body fluids or to determine its concentration. To prepare an immunoadsorbent for the purification of guinea pig IgG from serum or plasma.

Purification: Delipidated, heat inactivated, stable whole serum

Sterility: Heat-inactivated

Target Details

Target: IgG

Abstract: [IgG Products](#)

Target Type: Antibody

Application Details

Application Notes: In immunoelectrophoresis use 2 μ L serum or equivalent against 120 μ L antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ L antiserum in 3 mm diameter center well and 2 μ L serum samples (neat and serially diluted in 2 mm diameter peripheral wells).

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Reconstitute the lyophilized antiserum by adding 1 mL sterile distilled water. Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized antiserum by adding 1 mL sterile distilled water.

Buffer: Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added.

Preservative: Without preservative

Storage: RT, 4 °C, -20 °C

Storage Comment: The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C, prolonged storage at or below -20°C. Dilutions may be prepared by adding phosphate buffered

saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day.