

Datasheet for ABIN457748

Goat anti-Guinea Pig IgG1 (Fc Region) Antibody



()	ve	r\/i	Δ	۱۸/
\circ	V C	1 V		v v

Overview	
Quantity:	1 mL
Target:	lgG1
Binding Specificity:	Fc Region
Reactivity:	Guinea Pig
Host:	Goat
Clonality:	Polyclonal
Application:	Immunodiffusion (ID), Immunoelectrophoresis (IEP)
Product Details	
Immunogen:	Pools of purified normal IgG1 isolated from pooled guinea pig serum. Freund's complete
	adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Specificity:	The reactivity of the antiserum is directed to the Fc subunit of subclass IgG1.
Characteristics:	Precipitating polyclonal goat antiserum to guinea pig IgG1, subclass specific
	Total protein and IgG concentration in the antiserum are comparable to those of pooled normal
	goat serum. Antibody titre: Precipitin titre not less then 1:16 when tested against pooled normal
	guinea pig serum in agar block immun
Purification:	Delipidated, heat inactivated, lyophilized whole antiserum.
	Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies cross-
	reacting with other components of the immunoglobulin system or reacting with other serum
	proteins. Special attention is given to the removal of antibodies to common Ig/Fab. The use of

insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.

Target Details

Target:	lgG1
Abstract:	IgG1 Products
Target Type:	Antibody

Application Details

Application Notes:	In precipitating techniques as immunoelectrophoresis and radial immunodiffusion to identify
	the presence of IgG1 in guinea pig serum and other body fluids or to determine its concen-
	tration. To prepare an immunoadsorbent for the purification of guinea pig IgG1 from serum or
	plasma.
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Reconstitute the lyophilized antiserum by adding 1 mL sterile distilled water.
Buffer:	No foreign proteins added.
Preservative:	Without preservative
Handling Advice:	Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7. 2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4 °C, not refrozen, and preferably used the same day.
Storage:	4 °C/-20 °C
Storage Comment:	The lyophilized antiserum is shipped at ambient temperature and may be stored at +4 °C, prolonged storage at or below -24 °C.