

Datasheet for ABIN457769

## Rabbit anti-Guinea Pig IgG (Chain kappa), (Chain lambda), (Fab Region), (Free & Bound) Antibody



[Go to Product page](#)

### Overview

Quantity:	1 mL
Target:	IgG
Binding Specificity:	Chain kappa, Chain lambda, Fab Region, Free & Bound
Reactivity:	Guinea Pig
Host:	Rabbit
Clonality:	Polyclonal
Application:	Immunodiffusion (ID), Immunoelectrophoresis (IEP), Radial Immunodiffusion (RID)

### Product Details

Immunogen:	Purified Fab of normal guinea pig IgG. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Specificity:	The antiserum is reacting with the Fab subunit of intact IgG, IgA and IgM and other Ig classes of both light chain types, with their Fab or F(ab') <sub>2</sub> subunits and with free light chains of kappa and lambda type as tested in immunoelectrophoresis and double radial immunodiffusion
Characteristics:	<p>Precipitating polyclonal rabbit antiserum to guinea pig Fab of IgG</p> <p>Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit serum. Antibody titre: Precipitin titre not less than 1:32 when tested against normal guinea pig serum in agar block titration.</p>
Purification:	<p>Delipidated, heat inactivated lyophilized stable whole serum</p> <p>Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies</p>

## Product Details

---

reacting with other subunits of the immunoglobulin or reacting with other serum proteins. The use of insolubilized adsorption antigens prevents the presence of foreign protein or immune complexes in the antiserum.

## Target Details

---

Target: IgG

Abstract: [IgG Products](#)

Target Type: Antibody

## Application Details

---

Application Notes: In immunoelectrophoresis and double radial immunodiffusion to identify the presence of Fab, free or bound.

Restrictions: For Research Use only

## Handling

---

Format: Lyophilized

Reconstitution: It is reconstituted by adding 1 mL sterile distilled water.

Buffer: No foreign proteins added.

Preservative: Without preservative

Handling Advice: Working dilutions are prepared by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4 °C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.

Storage: 4 °C/-20 °C

Storage Comment: The lyophilized antiserum is shipped at ambient temperature and may be stored at +4 °C, prolonged storage at or below -24 °C.