

Datasheet for ABIN458069

**Goat anti-Human IgE (Fc Region) Antibody (TRITC)**[Go to Product page](#)

## Overview

Quantity:	1 mL
Target:	IgE
Binding Specificity:	Fc Region
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	TRITC
Application:	ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	Purified monoclonal IgE isolated from pooled human serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Specificity:	Specificity Reference Reagent. Tetramethylrhodamine isothiocyanate-conjugated IgG fraction of polyclonal Goat antiSerum to Human IgE, Fc specific.
Cross-Reactivity (Details):	Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this immunoconjugate has not been tested in detail.
Characteristics:	The reactivity of the antiserum is directed to the Fc subunit of the IgE molecule which expresses strict isotypic (class) specificity. It does not react with any non-Ig protein in human serum, as tested by immunoelectrophoresis and double radial immunodiffusion. Direct

## Product Details

---

immunofluorescence staining of cytoplasmic IgE of appropriately treated cell and tissue substrates, to demonstrate immunoglobulins or specific antibodies in cells and tissues, to identify circulating antibodies in serodiagnostic microbiology and autoimmune diseases. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.

Purification: Purified

## Target Details

---

Target: IgE

Abstract: [IgE Products](#)

## Application Details

---

Application Notes: ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.

Restrictions: For Research Use only

## Handling

---

Format: Lyophilized

Reconstitution: It is reconstituted by adding 1 mL sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20 °C. TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 mL sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20 °C.

Buffer: TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.

Preservative: Without preservative

Storage: RT, 4 °C, -20 °C

Storage Comment: The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C, prolonged storage at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile

phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.