

[Go to Product page](#)

Datasheet for ABIN458171

Sheep anti-Human IgD (Fc Region) Antibody (FITC)

Overview

| | |
|----------------------|---|
| Quantity: | 1 mL |
| Target: | IgD |
| Binding Specificity: | Fc Region |
| Reactivity: | Human |
| Host: | Sheep |
| Clonality: | Polyclonal |
| Conjugate: | FITC |
| Application: | ELISA, Immunofluorescence (IF), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC) |

Product Details

| | |
|-----------------------------|---|
| Immunogen: | Pools of purified homogenous IgD isolated from human serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Specificity: | Fluorescein isothiocyanate-conjugated IgG fraction of polyclonal Sheep antiSerum to HumanIgD Fc specific. |
| Cross-Reactivity (Details): | This immunoconjugate is not species-specific since inter-species cross-reactivity is a normal feature of antisera to immunoglobulins. Cross-reactivity of this conjugate has not been tested in detail. |
| Characteristics: | The reactivity of the antiserum is directed to the Fc subunit of the IgD molecule, which expresses strict (class) specificity. In immunoelectrophoresis and radial in immunodiffusion using various antiserum concentrations against human serum, a single precipitin line has been |

Product Details

obtained which shows a reaction of identity with the precipitin lines obtained with the purified IgD used as immunogens. It does not react with IgG including all subclasses, IgG/Fab fragments, IgM and IgA or any non-Ig protein in human serum, as tested by immunoelectrophoresis and double radial immunodiffusion. The antiserum also react with membrane-bound IgD in peripheral blood cells as tested by immunofluorescence microscopy. In immunocytochemical and immunohistochemical staining for the detection of IgD at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates, to identify and measure IgD in human serum or other body fluids. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.

Purification: Purified

Target Details

Target: IgD

Abstract: [IgD Products](#)

Target Type: Antibody

Application Details

Application Notes: ELISA,Immunocytochemistry,Immunohistochemistry (frozen),(In)direct immunofluorescence.

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: It is reconstituted by adding 1 mL sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20 °C.FITC-coupled purified hyperimmune sheep IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 mL sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20 °C.

Buffer: FITC-coupled purified hyperimmune sheep IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody

Handling

activity.

| | |
|---------------|----------------------|
| Preservative: | Without preservative |
|---------------|----------------------|

| | |
|----------|----------------|
| Storage: | RT,4 °C,-20 °C |
|----------|----------------|

| | |
|------------------|--|
| Storage Comment: | <p>The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C, prolonged storage at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.</p> |
|------------------|--|