



[Go to Product page](#)

Datasheet for ABIN458263

## Goat anti-Monkey IgA, IgG, IgM (Heavy & Light Chain) Antibody

### Overview

Quantity:	1 mL
Target:	IgA, IgG, IgM
Binding Specificity:	Heavy & Light Chain
Reactivity:	Monkey
Host:	Goat
Clonality:	Polyclonal
Application:	Immunocytochemistry (ICC), Immunodiffusion (ID), Immunoelectrophoresis (IEP), Radial Immunodiffusion (RID)

### Product Details

Immunogen:	Purified normal IgG, IgA and IgM fractions isolated from pooled rhesus monkey serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Specificity:	The reactivity of the antiserum is directed to the Fc and the Fab subunits of the major immunoglobulins.
Characteristics:	Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum. Antibody titre: Precipitin titre not less than 1:32 when tested against normal goat serum in agar block titration.
Purification:	Delipidated, heat inactivated, lyophilized stable whole serum Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies cross-reacting with other serum proteins. The use of insolubilized adsorption antigens prevents the

## Product Details

---

presence of excess adsorbent protein or immune complexes in the antiserum.

## Target Details

---

Target:	IgA, IgG, IgM
Alternative Name:	IgG + IgA + IgM ( <a href="#">IgA</a> , <a href="#">IgG</a> , <a href="#">IgM Products</a> )
Target Type:	Antibody

## Application Details

---

**Application Notes:** In immunoelectrophoresis and single and double radial immunodiffusion (Mancini, Ouchterlony) to identify and measure immunoglobulins, as secondary antibody to precipitate the immunoglobulin in normal monkey serum. This antiserum is not intended for use in non-precipitating antibody-binding or other highly sensitive assays. For such uses labelled and unlabelled cytochemical and immunoassay grade reagents meeting the necessary additional specificity and performance requirements are available.

**Restrictions:** For Research Use only

## Handling

---

Format:	Lyophilized
Reconstitution:	It is reconstituted by adding 1 mL sterile distilled water.
Buffer:	No foreign proteins added.
Preservative:	Without preservative
Handling Advice:	Working dilutions are prepared by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4 °C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.
Storage:	4 °C/-20 °C
Storage Comment:	The lyophilized antiserum is shipped at ambient temperature and may be stored at +4 °C, prolonged storage at or below -24 °C.