

Datasheet for ABIN458324

**Rabbit anti-Monkey IgG (Fab Region) Antibody - Preadsorbed**[Go to Product page](#)

## Overview

Quantity:	10 mg
Target:	IgG
Binding Specificity:	Fab Region
Reactivity:	Monkey
Host:	Rabbit
Clonality:	Polyclonal
Application:	ELISA, Immunohistochemistry (IHC), Immunocytochemistry (ICC)

## Product Details

Immunogen:	Purified Fab from normal IgG isolated from pooled Rhesus monkey serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Specificity:	The antiserum is reacting with the Fab subunit of intact IgG, IgA and IgM and other Ig classes of both light chain types, with their Fab or F(ab') <sub>2</sub> subunits and with free light chains of kappa and lambda type as tested in immunoelectrophoresis and double radial immunodiffusion
Characteristics:	Purified IgG fraction of polyclonal rabbit antiserum to monkey Fab of IgG Antibody titre: Precipitin titre not less than 1:64 when tested against normal monkey serum in agar block titration.
Purification:	Preadsorption: Immunoaffinity adsorbed using insolubilized antigens.

## Target Details

---

Target: IgG

Abstract: [IgG Products](#)

Target Type: Antibody

## Application Details

---

**Application Notes:** For indirect staining of fixed cell and tissue substrates, to demonstrate the intracellular presence of free or Ig-bound subunits of both kappa and lambda type. In general this kind of products is not recommended as direct or indirect screening reagents for immunoglobulin isotypes on the surface of membranes of vital lymphoid cells. The presence of activity to the common Fab subunit may result in the staining of Ig bound to Fc-receptors on non-lymphoid cells. Combinations of isotype-specific reagents should be used instead for this purpose. When applied in any cytochemical or histochemical procedure or solids phase coupling technique, the optimum concentration of the IgG preparation should always be established by titration. Typical working dilutions in histochemistry are usually between 1:50 and 1:500, in ELISA and comparable non-precipitating antibody-binding assays between 1:1000 and 1:5,000.

**Restrictions:** For Research Use only

## Handling

---

**Format:** Lyophilized

**Reconstitution:** It is reconstituted by adding 1 mL sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -24 °C.

**Concentration:** 10 mg/mL

**Buffer:** Purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2).

**Preservative:** Without preservative

**Handling Advice:** Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4 °C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.

**Storage:** 4 °C/-20 °C

## Handling

---

Storage Comment: The lyophilized IgG fraction is shipped at ambient temperature and may be stored at +4 °C, prolonged storage at or below -24 °C.