

Datasheet for ABIN458337

anti-Transferrin antibody



Overview

| Quantity: | 1 mL |
|------------------|--|
| Target: | Transferrin (TF) |
| Reactivity: | Mouse |
| Host: | Goat |
| Clonality: | Polyclonal |
| Conjugate: | This Transferrin antibody is un-conjugated |
| Application: | ELISA, Immunodiffusion (ID), Immunoelectrophoresis (IEP), Nephelometry (Neph), Radial Immunodiffusion (RID) |
| Product Details | |
| Immunogen: | Purified transferrin isolated from pooled mouse serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Isotype: | IgG |
| Specificity: | Inter-species cross-reactivity is a normal feature of antibodies to serum proteins, since homologous proteins of different species frequently share antigenic determinants. of this antiserum has not been tested in detail. |
| Characteristics: | Precipitating polyclonal goat antiserum to mouse transferrin |
| Purification: | Adsorption: Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies reacting with other serum proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum. |

Target Details

| Target: | Transferrin (TF) |
|---------------------|---|
| Alternative Name: | Transferrin (TF Products) |
| Background: | The reactivity of the antiserum is restricted to transferrin as tested in immunoelectro-phoresis and radial immunodiffusion. A characteristic single precipitin line is obtained with normal serum which shows a reaction of identity with the precipitin line with purified transferrin |
| Pathways: | Transition Metal Ion Homeostasis |
| Application Details | |
| Application Notes: | In precipitating techniques as immunoelectrophoresis and single and double radial immunodiffusion to identify the presence transferrin in mouse serum or other body fluids and to determine its concentration. This antiserum has not been tested in more sensitive assays as nephelometry, ELISA or use in immunohistology, but this does not exclude such use if proper controls are included. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Lyophilized |
| Concentration: | Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Antibody titre: Precipitin titre 1:64 when tested against pooled normal mouse serum in agar-block immunodiffusion titrati |
| Buffer: | Delipidated, heat inactivated, lyophilized, stable whole antiserum |
| Preservative: | Without preservative |
| Storage: | 4 °C/-20 °C |
| Storage Comment: | The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C, prolonged storage at or below -20°C. Reconstitute the lyophilized antiserum by add ing 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day. |