

Datasheet for ABIN458632

anti-Albumin antibody



Overview

Overview	
Quantity:	1 mL
Target:	Albumin (ALB)
Reactivity:	Rat
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This Albumin antibody is un-conjugated
Application:	ELISA, Immunoelectrophoresis (IEP), Immunodiffusion (ID), Radial Immunodiffusion (RID), Nephelometry (Neph)
Product Details	
Immunogen:	Highly purified albumin isolated from rat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Specificity:	Inter-species cross-reactivity is a normal feature of antibodies to mammalian serum proteins, since homologous proteins of different species frequently share antigenic determinants. The degree of cross-reactivity is also dependent on the concentrations of the reactants and the sensitivity of the assay arrangement. This antiserum fraction has been tested for cross-reactivity by double radial immunodiffusion against several species sera with the following results: cow - chicken - dog + goat - guinea pig ++ horse + human + mouse ++ monkey + rabbit
	± sheep - swine +

Product Details Purification:

Storage Comment:

Adsorption: Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies reacting with other serum proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.

Target:	Albumin (ALB)
Alternative Name:	Serum Albumin (ALB Products)
Background:	The defined antibody specificity is directed to albumin as tested against rat sera. In immuno-
	electrophoresis and double radial immunodiffusion (Ouchterlony), using various antiserum
	concentrations against appropriate concentrations of the immunogen, a single characteristic
	precipitin line is obtained which shows a reaction of identity with the precipitin lines obtained
	against rat serum and the purified albumin.
Pathways:	Lipid Metabolism
Application Details	
Application Notes:	As precipitating antiserum to identify or measure rat albumin by a variety of immunodiffusion
	techniques, including immunoelectrophoresis, single and double radial immunodiffusion
	(Mancini, Ouchterlony) and electroimmunodiffusion (Laurell). It has not been tested for use in
	nephelometry, ELISA or immunochemistry, but this does not exclude such use if proper
	controls are included.
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Concentration:	Total protein and IgG concentrations in the antiserum are comparable to those in pooled goat
	serum. No foreign proteins added. Antibody titre: Precipitin titre not less than 1:64 when tested
	against normal rat serum in agar block immunodiffusion titration
Buffer:	Delipidated, heat inactivated, lyophilized, stable whole antiserum
Preservative:	Without preservative
Storage:	4 °C/-20 °C

The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C,

prolonged storage at or below -20°C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not ref rozen, and preferably used the same day.