

## Datasheet for ABIN458888

## anti-CTSC antibody



| _ |       |             |    |    |             |     |
|---|-------|-------------|----|----|-------------|-----|
|   | V     | $\triangle$ | r۱ | /1 | $\triangle$ | Λ/  |
|   | ' V ' |             | ΙV |    |             | v v |

| Quantity:         | 1 mg   |  |
|-------------------|--|--|
| Target:           | CTSC   |  |
| Reactivity:       | Cow  |  |
| Host:             | Rabbit   |  |
| Clonality:        | Polyclonal   |  |
| Conjugate:        | This CTSC antibody is un-conjugated  |  |
| Application:      | Western Blotting (WB), ELISA, Immunofluorescence (IF)  |  |
| Product Details   |  |  |
| Immunogen:        | Cathepsin C isolated and purified from bovine spleen. Freund's complete adjuvant is used in the first step of the immunization procedure.  |  |
| Opposition        |  |  |
| Specificity:      | Cross-reactivities against enzymes of other sources may occur but have not been determined.  |  |
| Characteristics:  | Affinity purified antibodies of polyclonal rabbit antiserum to cathepsin C from bovine spleen  |  |
| Purification:     | Specific polyclonal antibody fractions were prepared by solid phase affinity chromatography.   |  |
| Target Details    |  |  |
| Target:           | CTSC   |  |
| Alternative Name: | Cathepsin C (CTSC Products)  |  |
| Background:       | The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: immunoelectrophoresis, cross-immunoelectrophoresis, single radial |  |
|                   | immunodiffusion (Ouchterlony), block titration, ELISA, immunoblotting and enzyme inhibition.   |  |

## **Application Details**

| Application Notes: | This product is intended for use in precipitating and non-precipitating antibody-binding assays (such as e.g., ELISA and Western blotting and immunofluorescence or histochemical techniques), to prepare an insoluble immuno-affinity adsorbent, for labelling with a marker of the customer's own choice.  |  |  |
|--------------------|--|--|--|
| Restrictions:      | For Research Use only  |  |  |
| Handling           |  |  |  |
| Format:            | Lyophilized  |  |  |
| Concentration:     | PAb concentration 1 mg/0.5ml. No foreign proteins added.   |  |  |
| Buffer:            | Purified hyperimmune rabbit IgG antibodies lyophilised from a solution in phosphate buffered saline (PBS, pH 7.2), stabilized with dextran.  |  |  |
| Preservative:      | Without preservative   |  |  |
| Storage:           | 4 °C/-20 °C  |  |  |
| Storage Comment:   | The lyophilised PAb fraction is shipped at ambient temperature and may be stored at +4°C, prolonged storage at or below -20°C. It is reconstituted by adding 0.5 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at a mbient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product. |  |  |