



[Go to Product page](#)

Datasheet for ABIN459084

## anti-Nucleoside Monophosphate Kinase antibody (Biotin)

### Overview

|              |   |
|--------------|---|
| Quantity:    | 1 mL  |
| Target:      | Nucleoside Monophosphate Kinase                                       |
| Reactivity:  | Cow   |
| Host:        | Rabbit  |
| Clonality:   | Polyclonal  |
| Conjugate:   | This Nucleoside Monophosphate Kinase antibody is conjugated to Biotin |
| Application: | Western Blotting (WB), Immunofluorescence (IF), ELISA                 |

### Product Details

|                  |  |
|------------------|--|
| Immunogen:       | Nucleoside monophosphate kinase isolated and purified from bovine liver. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Isotype:         | IgG  |
| Specificity:     | Cross-reactivities against enzymes of other sources may occur but have not been determined.  |
| Characteristics: | Biotin-conjugated IgG fraction of polyclonal rabbit antiserum to nucleoside mono-phosphate kinase from bovine liver  |
| Purification:    | The IgG (7S) fraction is prepared from the antiserum by ammonium sulphate precipitation and ion exchange chromatography.                                     |

### Target Details

|                   |   |
|-------------------|---|
| Target:           | Nucleoside Monophosphate Kinase   |
| Alternative Name: | Nucleoside Mono-phosphate Kinase ( <a href="#">Nucleoside Monophosphate Kinase Products</a> ) |

## Target Details

|             |   |
|-------------|---|
| Background: | The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: immunoelectrophoresis, cross-immunoelectrophoresis, single radial immunodiffusion (Ouchterlony), block titration, ELISA, immunoblotting and enzyme inhibition. |
|-------------|---|

## Application Details

|                    |   |
|--------------------|---|
| Application Notes: | This product is intended for use in precipitating and non-precipitating antibody-binding assays (such as e.g., ELISA and Western blotting and immuno-fluorescence or histochemical techniques). |
|--------------------|---|

|               |                       |
|---------------|-----------------------|
| Restrictions: | For Research Use only |
|---------------|-----------------------|

## Handling

|                  |  |
|------------------|--|
| Format:          | Lyophilized  |
| Concentration:   | IgG protein concentration 10 mg/ml. Biotin/IgG protein molar ratio (B/P) approximately 5.3. No foreign proteins added.   |
| Buffer:          | Biotin-coupled hyperimmune rabbit IgG lyophilised from a solution in phosphate buffered saline (PBS, pH 7.2).  |
| Preservative:    | Without preservative   |
| Storage:         | 4 °C/-20 °C  |
| Storage Comment: | The lyophilised conjugate is shipped at ambient temperature and may be stored at +4°C, prolonged storage at or below -20°C. It is reconstituted by adding 1.0 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at a mbient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used t he same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product. |