antibodies -online.com





anti-Phospholipase D antibody (Biotin)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

first step of the immunization procedure. Isotype: IgG Specificity: Cross-reactivities against enzymes of other sources may occur but have not been determined. Characteristics: Biotin-conjugated IgG fraction of polyclonal rabbit antiserum to phospholipase D from cabba	Overview		
Reactivity: Cabbage Host: Rabbit Clonality: Polyclonal Conjugate: This Phospholipase D antibody is conjugated to Biotin Application: Western Blotting (WB), Immunofluorescence (IF), ELISA Product Details Immunogen: Phospholipase D isolated and purified from cabbage. Freund's complete adjuvant is used infirst step of the immunization procedure. Isotype: IgG Specificity: Cross-reactivities against enzymes of other sources may occur but have not been determined that the procedure of the immunization of polyclonal rabbit antiserum to phospholipase D from cabbase. Purification: The IgG (7S) fraction is prepared from the antiserum by ammonium sulphate precipitation at ion exchange chromatography. Target Details Target: Phospholipase D (PLD)	Quantity:	1 mL	
Host: Rabbit Clonality: Polyclonal Conjugate: This Phospholipase D antibody is conjugated to Biotin Application: Western Blotting (WB), Immunofluorescence (IF), ELISA Product Details Immunogen: Phospholipase D isolated and purified from cabbage. Freund's complete adjuvant is used in first step of the immunization procedure. Isotype: IgG Specificity: Cross-reactivities against enzymes of other sources may occur but have not been determine. Characteristics: Biotin-conjugated IgG fraction of polyclonal rabbit antiserum to phospholipase D from cabbage. Purification: The IgG (7S) fraction is prepared from the antiserum by ammonium sulphate precipitation and ion exchange chromatography. Target Details Target: Phospholipase D (PLD)	Target:	Phospholipase D (PLD)	
Clonality: Polyclonal Conjugate: This Phospholipase D antibody is conjugated to Biotin Application: Western Blotting (WB), Immunofluorescence (IF), ELISA Product Details Immunogen: Phospholipase D isolated and purified from cabbage. Freund's complete adjuvant is used in first step of the immunization procedure. Isotype: IgG Specificity: Cross-reactivities against enzymes of other sources may occur but have not been determined that cateristics: Biotin-conjugated IgG fraction of polyclonal rabbit antiserum to phospholipase D from cabbase Purification: The IgG (7S) fraction is prepared from the antiserum by ammonium sulphate precipitation are ion exchange chromatography. Target Details Target: Phospholipase D (PLD)	Reactivity:	Cabbage	
Conjugate: This Phospholipase D antibody is conjugated to Biotin Application: Western Blotting (WB), Immunofluorescence (IF), ELISA Product Details Immunogen: Phospholipase D isolated and purified from cabbage. Freund's complete adjuvant is used infirst step of the immunization procedure. Isotype: IgG Specificity: Cross-reactivities against enzymes of other sources may occur but have not been determined that the conjugated IgG fraction of polyclonal rabbit antiserum to phospholipase D from cabbase Purification: The IgG (7S) fraction is prepared from the antiserum by ammonium sulphate precipitation and ion exchange chromatography. Target Details Target: Phospholipase D (PLD)	Host:	Rabbit	
Application: Western Blotting (WB), Immunofluorescence (IF), ELISA Product Details Immunogen: Phospholipase D isolated and purified from cabbage. Freund's complete adjuvant is used in first step of the immunization procedure. Isotype: IgG Specificity: Cross-reactivities against enzymes of other sources may occur but have not been determined. Characteristics: Biotin-conjugated IgG fraction of polyclonal rabbit antiserum to phospholipase D from cabbase. Purification: The IgG (7S) fraction is prepared from the antiserum by ammonium sulphate precipitation and ion exchange chromatography. Target Details Target: Phospholipase D (PLD)	Clonality:	Polyclonal	
Product Details Immunogen: Phospholipase D isolated and purified from cabbage. Freund's complete adjuvant is used in first step of the immunization procedure. Isotype: IgG Specificity: Cross-reactivities against enzymes of other sources may occur but have not been determined. Characteristics: Biotin-conjugated IgG fraction of polyclonal rabbit antiserum to phospholipase D from cabbase. Purification: The IgG (7s) fraction is prepared from the antiserum by ammonium sulphate precipitation and ion exchange chromatography. Target Details Target: Phospholipase D (PLD)	Conjugate:	This Phospholipase D antibody is conjugated to Biotin	
Immunogen: Phospholipase D isolated and purified from cabbage. Freund's complete adjuvant is used in first step of the immunization procedure. Isotype: IgG Specificity: Cross-reactivities against enzymes of other sources may occur but have not been determined the conjugated IgG fraction of polyclonal rabbit antiserum to phospholipase D from cabbate Purification: The IgG (7S) fraction is prepared from the antiserum by ammonium sulphate precipitation and ion exchange chromatography. Target Details Target: Phospholipase D (PLD)	Application:	Western Blotting (WB), Immunofluorescence (IF), ELISA	
first step of the immunization procedure. Isotype: IgG Specificity: Cross-reactivities against enzymes of other sources may occur but have not been determined. Characteristics: Biotin-conjugated IgG fraction of polyclonal rabbit antiserum to phospholipase D from cabbate Purification: The IgG (7S) fraction is prepared from the antiserum by ammonium sulphate precipitation are ion exchange chromatography. Target Details Target: Phospholipase D (PLD)	Product Details		
Isotype: IgG Specificity: Cross-reactivities against enzymes of other sources may occur but have not been determined. Characteristics: Biotin-conjugated IgG fraction of polyclonal rabbit antiserum to phospholipase D from cabbate. Purification: The IgG (7S) fraction is prepared from the antiserum by ammonium sulphate precipitation at ion exchange chromatography. Target Details Target: Phospholipase D (PLD)	Immunogen:	Phospholipase D isolated and purified from cabbage. Freund's complete adjuvant is used in the	
Specificity: Cross-reactivities against enzymes of other sources may occur but have not been determined. Characteristics: Biotin-conjugated IgG fraction of polyclonal rabbit antiserum to phospholipase D from cabbate. Purification: The IgG (7S) fraction is prepared from the antiserum by ammonium sulphate precipitation and ion exchange chromatography. Target Details Target: Phospholipase D (PLD)		first step of the immunization procedure.	
Characteristics: Biotin-conjugated IgG fraction of polyclonal rabbit antiserum to phospholipase D from cabba Purification: The IgG (7S) fraction is prepared from the antiserum by ammonium sulphate precipitation are ion exchange chromatography. Target Details Target: Phospholipase D (PLD)	Isotype:	IgG	
Purification: The IgG (7S) fraction is prepared from the antiserum by ammonium sulphate precipitation are ion exchange chromatography. Target Details Target: Phospholipase D (PLD)	Specificity:	Cross-reactivities against enzymes of other sources may occur but have not been determined.	
Target Details Target: Phospholipase D (PLD)	Characteristics:	Biotin-conjugated IgG fraction of polyclonal rabbit antiserum to phospholipase D from cabbage	
Target Details Target: Phospholipase D (PLD)	Purification:	The IgG (7S) fraction is prepared from the antiserum by ammonium sulphate precipitation and	
Target: Phospholipase D (PLD)		ion exchange chromatography.	
	Target Details		
Alternative Name: Phospholipase D (PLD Products)	Target:	Phospholipase D (PLD)	
	Alternative Name:	Phospholipase D (PLD Products)	

Target Details	
Background:	The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: immunoelectrophoresis, cross-immunoelectrophoresis, single radial immunodiffusion (Ouchterlony), block titration, ELISA, immunoblotting and enzyme inhibition.
Pathways:	Response to Water Deprivation, ER-Nucleus Signaling
Application Details	
Application Notes:	This product is intended for use in precipitating and non-precipitating antibody-binding assays (such as e.g., ELISA and Western blotting and immunofluorescence or histochemical techniques).
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Concentration:	IgG protein concentration 10 mg/ml. Biotin/IgG protein molar ratio (B/P) approximately 5.5. No foreign proteins added.
Buffer:	Biotin-coupled hyperimmune rabbit IgG lyophilised from a solution in phosphate buffered saline (PBS, pH 7.2).
Preservative:	Without preservative
Storage:	4 °C/-20 °C
Storage Comment:	The lyophilised conjugate is shipped at ambient temperature and may be stored at +4°C, prolonged storage at or below -20°C. It is reconstituted by adding 1.0 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at a mbient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon

product.

storage, this should be removed by centrifugation. It will not affect the performance of the