



[Go to Product page](#)

Datasheet for ABIN459210  
**anti-Glycerol Dehydrogenase antibody**

### Overview

Quantity:	10 mg
Target:	Glycerol Dehydrogenase
Reactivity:	Klebsiella aerogenes
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Glycerol Dehydrogenase antibody is un-conjugated
Application:	Immunofluorescence (IF), Western Blotting (WB), ELISA

### Product Details

Immunogen:	Glycerol dehydrogenase isolated and purified from Enterobacter aerogenes. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Specificity:	Cross-reactivities against enzymes of other sources may occur but have not been determined.
Characteristics:	IgG fraction of polyclonal rabbit antiserum to glycerol dehydrogenase from Enterobacter aerogenes
Purification:	The IgG (7S) fraction is prepared from the antiserum by ammonium sulphate precipitation and ion exchange chromatography.

### Target Details

Target:	Glycerol Dehydrogenase
Abstract:	<a href="#">Glycerol Dehydrogenase Products</a>

## Target Details

---

**Background:** The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: immunoelectrophoresis, cross-immunoelectrophoresis, single radial immunodiffusion (Ouchterlony), block titration, ELISA, immunoblotting and enzyme inhibition.

## Application Details

---

**Application Notes:** This product is intended for use in precipitating and non-precipitating antibody-binding assays (such as e.g., ELISA and Western blotting and immunofluorescence or histochemical techniques), to prepare an insoluble immuno-affinity adsorbent, for labelling with a marker of the customer's own choice.

**Restrictions:** For Research Use only

## Handling

---

**Format:** Lyophilized

**Concentration:** IgG protein concentration 10 mg/ml. No foreign proteins added.

**Buffer:** Purified hyperimmune rabbit IgG lyophilised from a solution in phosphate buffered saline (PBS, pH 7.2).

**Preservative:** Without preservative

**Storage:** 4 °C/-20 °C

**Storage Comment:** The lyophilised IgG fraction is shipped at ambient temperature and may be stored at +4°C, prolonged storage at or below -20°C. It is reconstituted by adding 1.0 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.