antibodies -online.com





Datasheet for ABIN459256

anti-PLA2G1B antibody



Overview

Quantity:	10 mg
Target:	PLA2G1B
Reactivity:	Bee
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PLA2G1B antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF)
Product Details	
Immunogen:	Phospholipase A 2 isolated and purified from bee venom. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Specificity:	Cross-reactivities against enzymes of other sources may occur but have not been determined.
Characteristics:	IgG fraction of polyclonal rabbit antiserum to phospholipase A 2 from bee venom
Purification:	The IgG (7S) fraction is prepared from the antiserum by ammonium sulphate precipitation and ion exchange chromatography.
Target Details	
Target:	PLA2G1B
Alternative Name:	Phospholipase A 2 (PLA2G1B Products)

Target Details Background: The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: immunoelectrophoresis, cross-immunoelectrophoresis, single radial immunodiffusion (Ouchterlony), block titration, ELISA, immunoblotting and enzyme inhibition. Inositol Metabolic Process, VEGF Signaling Pathways: **Application Details** This product is intended for use in precipitating and non-precipitating antibody-binding assays Application Notes: (such as e.g., ELISA and Western blotting and immunofluorescence or histochemical techniques), to prepare an insoluble immuno-affinity adsorbent, for labelling with a marker of the customer's own choice. Restrictions: For Research Use only Handling Format: Lyophilized Concentration: IgG protein concentration 10 mg/ml. No foreign proteins added. Buffer: Purified hyperimmune rabbit IgG lyophilised from a solution in phosphate buffered saline (PBS, pH 7.2). Preservative: Without preservative Storage: 4 °C/-20 °C Storage Comment: The lyophilised IgG fraction is shipped at ambient temperature and may be stored at +4°C, prolonged storage at or below -20°C. It is reconstituted by adding 1.0 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or

below -20°C. Prior to use, an aliquot is thawed slowly at a mbient temperature, spun down