



[Go to Product page](#)

Datasheet for ABIN462493

## anti-Akirin 1 antibody (N-Term)

### 3 Images

#### Overview

Quantity:	0.05 mg
Target:	Akirin 1 (AKIRIN1)
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Akirin 1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (IHC)

#### Product Details

Brand:	IHC-plus™
Isotype:	IgG
Purification:	Immunoaffinity purified

#### Target Details

Target:	Akirin 1 (AKIRIN1)
Alternative Name:	AKIRIN1 ( <a href="#">AKIRIN1 Products</a> )
Background:	Name/Gene ID: AKIRIN1

Synonyms: AKIRIN1, Akirin 1, Akirin-1, RP11-781D11.2, STRF2, C1orf108

## Target Details

---

Gene ID: 79647

UniProt: [Q9H9L7](#)

## Application Details

---

Application Notes: Approved: IHC, IHC-P (5 µg/mL), WB (1 - 2 µg/mL)

Usage: Akirin1 antibody can be used for detection of Akirin1 by Western blot at 1-2 µg/mL

Comment: Target Species of Antibody: Human

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: Lot specific

Buffer: PBS, 0.02 % sodium azide

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling Advice: avoid freeze thaw cycles. Store undiluted.

Storage: 4 °C, -20 °C


Storage Comment: Store at 4°C for 3 months and -20°C, stable for up to 1 year. Avoid repeated freeze-thaw cycles.

Expiry Date: 12 months




**Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections)**

**Image 1.** Human Spleen: Formalin-Fixed, Paraffin-Embedded (FFPE)



**Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections)**

**Image 2.** Human Placenta: Formalin-Fixed, Paraffin-Embedded (FFPE)



**Immunohistochemistry**

**Image 3.** Anti-AKIRIN1 antibody IHC of human placenta. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 5 ug/ml.