antibodies - online.com







anti-FGF7 antibody (Biotin)

Images



| \sim | | | | |
|--------|------|-----|-----|-----|
| | 11/6 | ٦r١ | /10 | ۱۸. |

| Quantity: | 50 μg |
|-------------------|---|
| Target: | FGF7 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This FGF7 antibody is conjugated to Biotin |
| Application: | Western Blotting (WB), ELISA, Neutralization (Neut) |
| Product Details | |
| Immunogen: | E.coli derived recombinant Human KGF. |
| | Type of Immunogen: Cell extract |
| Specificity: | Human KGF |
| Purification: | Immunoaffinity purified |
| Target Details | |
| Target: | FGF7 |
| Alternative Name: | FGF7 / KGF (FGF7 Products) |
| Background: | Name/Gene ID: FGF7 |
| | Family: HBGF |

Target Details

| | Synonyms: FGF7, FGF-7, Fibroblast growth factor- 7, Keratinocyte growth factor, Fibroblast growth factor 7, HBGF-7, KGF |
|-----------|---|
| Gene ID: | 2252 |
| UniProt: | P21781 |
| Pathways: | RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway |

Application Details

Application Notes:

Approved: ELISA (1 µg/mL), Neut, WB (0.1 - 0.2 µg/mL)

Usage: The applications listed have been tested for the unconjugated form of this product. Other forms have not been tested. ELISA: To detect hKGF by direct ELISA (using 100 μ L/well antibody solution) a concentration of 0.25-1 μ g/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2-0.4 ng/well of recombinant hKGF. To detect hKGF by sandwich ELISA (using 100 μ L/well antibody solution) a concentration of 0.25-1 μ g/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with Polyclonal Anti-Human KGF (LS-C104693) as a capture antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant hKGF. Western Blot: To detect hKGF by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 μ g/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hKGF is 1.5-3 ng/lane, under either reducing or non-reducing conditions.

Comment:

Target Species of Antibody: Human

Restrictions:

For Research Use only

Handling

| Format: | Lyophilized |
|------------------|----------------------------------|
| Reconstitution: | Sterile PBS containing 0.1 % BSA |
| Concentration: | Lot specific |
| Buffer: | Lyophilized from PBS, pH 7.2 |
| Handling Advice: | Avoid freeze/thaw cycles. |

Handling

| Storage: | 4 °C,-20 °C |
|------------------|---|
| Storage Comment: | Store Lyophilized at room temperature up to 1 month Reconstituted for up to 2 weeks at 2-8°C. Aliquot and freeze at -20°C for long term storage. |
| | Avoid freeze/thaw cycles. |

Images




Image 1.




Image 2.