



[Go to Product page](#)

Datasheet for ABIN4880057

anti-CRISPR-Cas9 (N-Term) antibody

2 Validations

Overview

Quantity:	0.1 mL
Target:	CRISPR-Cas9
Binding Specificity:	N-Term
Reactivity:	Streptococcus pyogenes
Host:	Mouse
Clonality:	Monoclonal
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Immunogen:	This antibody was raised against a recombinant protein within the N-terminal region of Streptococcus pyogene Cas9
Clone:	7A9-3A3
Isotype:	IgG1 kappa
Specificity:	This antibody should recognize Cas9 and dCas9 based on the antigen design
Purification:	Protein A

Target Details

Target:	CRISPR-Cas9
Alternative Name:	CRISPR/Cas9

Target Details

Gene ID: 901176

Application Details

Application Notes: Optimal antibody dilution should be determined by titration, however as a guideline try at,IB 0.1-1 µg/mL

Comment: Myeloma, fusion partners: X63.Ag8-653

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 mg/mL

Buffer: Purified antibody (from supernatant) containing PBS + 0.09 % sodium azide

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.



Successfully validated (Western Blotting (WB))

by [Group Kleinlogel](#), Department of Physiology, University of Bern, Bern, Switzerland

Report Number: 104239

Date: May 28 2020

Target:	SpCas9
Lot Number:	1060814
Method validated:	Western Blotting (WB)
Positive Control:	HEK293 cells transfected with a plasmid expressing SpCas9-T2A-eGFP
Negative Control:	Non-transfected HEK293 cells
Notes:	Passed. ABIN4880057 specifically recognizes the Cas9 protein in HEK293 cells.
Primary Antibody:	ABIN4880057
Secondary Antibody:	HRP-conjugated goat anti-mouse antibodies (Jackson Immuno Research, 115-035-146)
Protocol:	<ul style="list-style-type: none">• Grow human embryonic kidney (HEK293) cells (ECACC, 85120602) in DMEM (Sigma, D5671, lot RNB68272) supplemented with 10% Fetal calf Serum (Seraglob, S70500, lot 208/203142), L-Alanyl-L-Glutamine (Merck, Cat. K0302), and Penicillin-Streptomycin (Sigma, P4333, lot 058M4857V) at 37 °C and 5% CO₂.• Plate 0.5x10⁶ cells/ml in 2 ml/well of cells in a 6 well plate.• Grow cells for 24 h at 37 °C and 5% CO₂.• Transfect cells with 2 µg/well of a plasmid expressing human codon optimized SpCas9-T2A-eGFP under the control of the ubiquitous promoter CMV using the calcium phosphate method. For each well, prepare 150 µl 0.25 M CaCl₂, add DNA, add 150 µl HBS, incubate for 3min at RT, and add the whole mix to the cells.• Change the media after 4-6 h.• Grow cells for 72 h at 37 °C and 5% CO₂.• Harvest cells with PBS and lyse them on ice for 30min with 50µl/well of RIPA buffer (25 mM TrisHCl pH7-8, 150 mM NaCl, 0.1% SDS, 0.5% sodium deoxycholate, 1% Triton X-100 or NP-40).• Complete the lysis with a freeze/taw cycle.• Determine total protein content of the lysates using Pierce BCA protein assay (Thermo Fisher, 23221).• Denature 25 µg of total protein for 5min at 95 °C in 20 µl Laemmli SDS sample buffer and subsequently separate them on a denaturing 4-20% Mini-PROTEAN TGX Stain-Free Gel (Bio-Rad, 456-8094) for 20 min at 100 V and 2 h at 130 V.• Transfer proteins onto Immobilon-P transfer membrane (Immobilion, IPVH00010) with a


Western blotting system for 75 min at 100 V.

- Block the membrane with TBST 5% milk for 1 h at RT.
- Incubate with primary
 - mouse anti-cas9 antibody (antibodies-online, ABIN4880057, batch n: 1060814) diluted 1:1,000 in TBST 5% milk overnight at 4 °C.
 - mouse anti-2A peptide antibody (antibodies-online, ABIN5074774, lot A-12) diluted 1:1,000 in TBST 5% milk overnight at 4 °C.
 - mouse anti-GAPDH antibody (Fitzgerald, 10R-G09a, lot 2417) diluted 1:40,000 in TBST 5% milk overnight at 4 °C.
- Wash membrane 3x for 5 min with TBST buffer.
- Incubate with secondary HRP-conjugated goat anti-mouse antibodies (Jackson Immuno Research, 115-035-146) diluted 1:3,000 in TBST 5% milk for 1 h at RT.
- Wash membrane 3x for 10 min with TBST buffer.
- Reveal protein bands on a ChemiDoc MP imaging system (Bio-Rad, 17001402) using Westar Sun ECL Substrate (Cyanagen, XLS063).

Experimental Notes:

- The insertion of a 2A peptide between 2 coding sequences leads to the translation of a unique protein which is then spliced in two proteins by the cell machinery. In this process, the 2A peptide stays bound to the C-terminus of the first protein and it can be used as a tag. In this experiment, the T2A peptide is bound to the Cas9 protein. As a consequence, the use of the T2A peptide or the Cas9 antibody should detect a band at the same molecular weight.
- The Cas9 antibody ABIN4880057 reveals a protein of the expected molecular weight of antigen in lysates of HEK293 cells. The protein bands are only visible in the positives but not the negative controls. Importantly, in the Cas9 blot, the intensity of the band appears stronger than in the T2A-peptide blot suggesting that the Cas9 antibody is stronger than the T2A peptide antibody.

Image for Validation report #104239



Validation image no. 1 for anti-CRISPR-Cas9 (N-Term) antibody (ABIN4880057)

Western blot analysis of cell lysates from HEK293 cells transfected (pos) or not transfected (neg) with a Cas9-T2A-eGFP expression plasmid. Staining was performed using anti-T2Apeptide (top-left), ABIN4880057 (top-right), or an anti-GAPDH loading control (bottom) antibody.



Successfully validated (Immunofluorescence (IF))

by [Group Kleinlogel](#), Department of Physiology, University of Bern, Bern, Switzerland

Report Number: 104240

Date: Jul 03 2020


Target:	SpCas9
Lot Number:	1060814
Method validated:	Immunofluorescence (IF)
Positive Control:	HEK cells transfected with a plasmid expressing SaCas9-T2A-eGFP anti-T2A antibody
Negative Control:	Non-transfected HEK293 cells
Notes:	Passed. ABIN4880057 specifically recognizes SaCas9 in HEK cells expressing SaCas9-T2A-eGFP.
Primary Antibody:	ABIN4880057
Secondary Antibody:	goat anti-mouse Cy3 (Invitrogen, A10521)
Protocol:	<ul style="list-style-type: none">• Grow human embryonic kidney HEK293 cells (ECACC, 85120602) in DMEM (Sigma, D5671, lot RNB68272) supplemented with 10% Fetal calf Serum (Seraglob, S70500, lot 208/203142), L-Alanyl-L-Glutamine (Merck, Cat. K0302), and Penicillin-Streptomycin (Sigma, P4333, lot 058M4857V) at 37 °C and 5% CO₂.• The day before plating, place a 15mm round glass/well in 24 well plate and incubate overnight with poly L-ornitine (Sigma, P4957).• Wash with PBS.• Plate 2x10⁵ cells/well in 500 µl in a 24 well plate with round glasses.• Grow cells for 24 h at 37 °C and 5% CO₂.• Transfect cells with 0.2 µg/well of a plasmid expressing human SpCas9-T2A-eGFP under the control of the ubiquitous promoter CMV using the calcium phosphate method. For each well, prepare 30 µl 0.25 M CaCl₂, add DNA, add 30 µl HBS, incubate for 3 min at RT, and add the whole mix to the cells.• Change the media after 4-6 h.• Grow cells for 72 h at 37 °C and 5% CO₂.• Remove the media and wash once with PBS.• Add 250 µL of 4% PFA and incubate for 7 min.• Wash 3 times with PBS.• Add 250 µL of 0.1 M glycine in PBS to block unreacted aldehydes.• Remove glycine and add blocking solution (5% goat serum + 1% BSA Serum in 0.3% Triton-

- X100 in 1 x TBS) for 60 min at RT.
- Dilute primary
 - mouse anti-Cas9 antibody (antibodies-online, ABIN5074774, Batch N: 1060814) 1:250 or
 - mouse anti-2A peptide antibody (antibodies-online, ABIN5074774, lot A-12) 1:250 in blocking solution.
 - Add 250 µL/well of primary antibody solution.
 - Cover the plate with aluminum foil and Incubate overnight at 4 °C in the dark in a shaking plate.
 - Wash wells 3 times with PBS.
 - Dilute secondary goat anti-mouse Cy3 (Invitrogen, A10521) 1:400 and DAPI (0.1 ug/ml) in blocking solution.
 - Add 250 µL/well of secondary antibody solution.
 - Wash wells 3 times with PBS.
 - Place the round glasses in microscope slides.
 - Add fluorescence mounting medium (DAKO, S3023, lot 10115314) and cover with coverslips.
 - Seal the edges of the coverslips with a clear nail polish.
 - Let the nail polish dry and store samples at 4 °C in dark.
 - Acquire images with Axiovert 200M with a 40X objective.

Experimental Notes:

- The insertion of a 2A peptide between 2 coding sequences leads to the translation of a unique protein which is then spliced in two proteins by the cell machinery. In this process, the 2A peptide stays bound to the C-terminus of the first protein and it can be used as a tag. In this experiment, the T2A peptide is bound to the SaCas9 protein and it can be used as a positive control. Moreover we expect that cells expressing eGFP should express the saCas9 protein as well.
- The SaCas9 antibody ABIN6972659 specifically labels the targeted antigen in HEK293 cells transfected with a plasmid encoding SaCas9-T2A-eGFP. Importantly, cells are always colabeled with eGFP and the antibody.
- As a control, the T2A antibody shows a similar pattern.

Image for Validation report #104240



Validation image no. 1 for anti-CRISPR-Cas9 (N-Term) antibody (ABIN4880057)

HEK-293 cells transfected with a plasmid expressing SpCas9-T2A-eGFP and stained with ABIN4880057 and a Cy3 conjugated anti-mouse antibody (first row), an anti-2A antibody and a Cy3 conjugated anti-mouse antibody (second row). Staining of transfected cells with the secondary antibody alone (third row) or non-transfected

cells with ABIN4880057 or the Cy3-conjugated secondary antibody (forth row) or the the 2A-antibody and Cy3 secondary (fifth row) served as negative controls.