

Datasheet for ABIN4881312
NUCB2 ELISA Kit[Go to Product page](#)

2 Publications

Overview

Quantity:	96 tests
Target:	NUCB2 (NES1)
Reactivity:	Human, Rat, Mouse
Method Type:	Competition ELISA
Detection Range:	0.1-1.000 ng/mL
Minimum Detection Limit:	0.1 ng/mL
Application:	ELISA

Product Details

Purpose:	Human/Mouse/Rat Nesfatin EIA Kit optimized for serum and cell culture supernatants. Competition-based ELISA on a 96-well strip plate.
Sample Type:	Cell Culture Supernatant, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This kit detects human, mouse and rat 82aa Nesfatin protein. No other active isoforms have been reported.
Sensitivity:	400 pg/mL
Characteristics:	<ul style="list-style-type: none">• Strip plates and additional reagents allow for use in multiple experiments• Quantitative protein detection• Establishes normal range

Product Details

- The best products for confirmation of antibody array data

Components:

- Pre-Coated 96-well Strip Microplate
- Wash Buffer
- Standard Peptide
- Assay Diluent(s)
- Biotinylated Peptide
- HRP-Streptavidin
- TMB One-Step Substrate
- Stop Solution
- Assay Diagram
- Positive Control Sample
- Capture Antibody
- User Manual

Material not included:

- Distilled or deionized water
- Precision pipettes to deliver 2 µL to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- 100 mL and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- Orbital shaker
- Aluminum foil
- Saran Wrap
- Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- SigmaPlot software (or other software that can perform four-parameter logistic regression models)

Target Details

Target:	NUCB2 (NES1)
Alternative Name:	Nesfatin (NES1 Products)
Gene ID:	4925
UniProt:	P80303

Application Details

Application Notes:	Recommended Dilution for serum and plasma samplesHuman: 2X / Mouse: 2X / Rat: 2X
Sample Volume:	100 µL
Plate:	Pre-coated

Application Details

Protocol:	<ol style="list-style-type: none">1. Prepare all reagents, samples and standards as instructed.2. Add 100 µL detection antibody to each well.3. Incubate 1.5 h at RT or O/N at 4 °C.4. Add 100 µL standard or sample to each well.5. Incubate 2.5 h at RT.6. Add 100 µL prepared streptavidin solution.7. Incubate 45 min at RT.8. Add 100 µL TMB One-Step Substrate Reagent to each well.9. Incubate 30 min at RT.10. Add 50 µL Stop Solution to each well.11. Read plate at 450 nm immediately.
-----------	---

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Storage:	-20 °C
Storage Comment:	Standard, biotinylated peptide, and positive control should be stored at -20°C after arrival. Avoid multiple freeze-thaws. The remaining kit components may be stored at 4°C. Opened microplate wells and antibody (Item N) may be stored for up to 1 month at 2° to 8°C. Return unused wells to the pouch containing desiccant pack and reseal along entire edge.
Expiry Date:	6 months

Publications

Product cited in:	<p>Serin, Bakacak, Ercan, Köstü, Avci, Arıkan, Kıran: "The evaluation of Nesfatin-1 levels in patients with and without intrauterine growth restriction." in: The journal of maternal-fetal & neonatal medicine, Vol. 29, Issue 9, pp. 1409-13, (2016) (PubMed).</p> <p>Abaci, Catli, Anik, Kume, Bober: "The relation of serum nesfatin-1 level with metabolic and clinical parameters in obese and healthy children." in: Pediatric diabetes, Vol. 14, Issue 3, pp. 189-95, (2013) (PubMed).</p>
-------------------	--