antibodies .- online.com





Datasheet for ABIN4881336

Urocortin ELISA Kit



Overview

Quantity:	96 tests
Target:	Urocortin (UCN)
Reactivity:	Mouse, Human, Rat
Method Type:	Competition ELISA
Detection Range:	0.1-1.000 ng/mL
Minimum Detection Limit:	0.1 ng/mL
Application:	ELISA
Product Details	
Purpose:	Human/Mouse/Rat Urocortin EIA Kit optimized for serum. Competition-based ELISA on a 96-well strip plate.
Sample Type:	Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This EIA kit is designed to detect human, mouse, and rat active Urocortin.
Sensitivity:	1.2 ng/mL
Characteristics:	 Strip plates and additional reagents allow for use in multiple experiments Quantitative protein detection Establishes normal range The best products for confirmation of antibody array data

Product Details

Components:

- · Pre-Coated 96-well Strip Microplate
- · Wash Buffer
- · Standard Peptide
- Assay Diluent(s)
- · Biotinylated Peptide
- · HRP-Streptavidin
- · TMB One-Step Substrate
- · Stop Solution
- · Assay Diagram
- · Positive Control Sample
- · Capture Antibody
- User Manual

Material not included:

- Distilled or deionized water
- Precision pipettes to deliver 2 µL to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- 100 mL and 1 liter graduated cylinders
- · Tubes to prepare standard and sample dilutions
- · Orbital shaker
- · Aluminum foil
- · Saran Wrap
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- SigmaPlot software (or other software that can perform four-parameter logistic regression models)

Target Details

Target:	Urocortin (UCN)
Alternative Name:	Urocortin (UCN Products)
Gene ID:	7349
UniProt:	P55089
Pathways:	Positive Regulation of Peptide Hormone Secretion, Hormone Activity, Negative Regulation of Hormone Secretion, cAMP Metabolic Process, Regulation of Cell Size, Feeding Behaviour

Application Details

Sample Volume:	100 μL
Plate:	Pre-coated

Application Details

Protocol:	1. Prepare all reagents, samples and standards as instructed.
	2. Add 100 µL detection antibody to each well.
	3. Incubate 1.5 h at RT or O/N at 4 °C.
	4. Add 100 μL standard or sample to each well.
	5. Incubate 2.5 h at RT.
	6. Add 100 μL prepared streptavidin solution.
	7. Incubate 45 min at RT.
	8. Add 100 µL TMB One-Step Substrate Reagent to each well.
	9. Incubate 30 min at RT.
	10. Add 50 μL Stop Solution to each well.
	11. Read plate at 450 nm immediately.
Restrictions:	For Research Use only
Handling	
Storage:	-20 °C
Storage Comment:	Standard, Biotinylated Urocortin peptide, and Positive Control should be stored at -20°C after
	arrival. Avoid multiple freeze-thaws. The remaining kit components may be stored at 4°C.
	Opened Microplate Wells and antibody (Item N) may be stored for up to 1 month at 2° to 8°C.
	Return unused wells to the pouch containing desiccant pack and reseal along entire edge.
Expiry Date:	6 months