antibodies -online.com





Datasheet for ABIN4881949

BCL10 ELISA Kit



Overview

Quantity:	96 tests
Target:	BCL10
Reactivity:	Human
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details	
Purpose:	Custom Human BCL-10 ELISA Kit.
Sample Type:	Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Lysate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA antibody pair recognizes Human Cochlin.
Characteristics:	 Strip plates and additional reagents allow for use in multiple experiments Quantitative protein detection Establishes normal range The best products for confirmation of antibody array data
Components:	 Pre-Coated 96-well Strip Microplate Wash Buffer Stop Solution Assay Diluent(s) Lyophilized Standard Biotinylated Detection Antibody

Product Details

- · Streptavidin-Conjugated HRP
- TMB One-Step Substrate

Material not included:

- · Distilled or deionized water
- Precision pipettes to deliver 2 µL to 1 µL volumes
- Adjustable 1-25 µL pipettes for reagent preparation
- 100 µL and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- Log-log graph paper or computer and software for ELISA data analysis

Target Details

Target:	BCL10
Alternative Name:	BCL-10 (BCL10 Products)
Gene ID:	8915
UniProt:	095999
Pathways:	TCR Signaling, Fc-epsilon Receptor Signaling Pathway, Activation of Innate immune Response, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, Tube Formation, Positive Regulation of Endopeptidase Activity, BCR Signaling, Ubiquitin Proteasome Pathway, S100 Proteins

Application Details

Sample Volume:	100 μL
Plate:	Pre-coated
Protocol:	 Prepare all reagents, samples and standards as instructed in the manual. Add 100 μL of standard or sample to each well. Incubate 2.5 h at RT or O/N at 4 °C. Add 100 μL of prepared biotin antibody to each well. Incubate 1 h at RT. Add 100 μL of prepared Streptavidin solution to each well. Incubate 45 min at RT. Add 100 μL of TMB One-Step Substrate Reagent to each well. Incubate 30 min at RT. Add 50 μL of Stop Solution to each well.
	10. Add 50 μL of Stop Solution to each well.11. Read at 450 nm immediately.

Application Details

Restrictions:	For Research Use only
Handling	
Storage:	-20 °C
Storage Comment:	The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it is recommended to store at -80°C.
Expiry Date:	6 months