antibodies -online.com





Datasheet for ABIN4882158

CD28 ELISA Kit



Overview

| Quantity: | 96 tests |
|--------------|----------------|
| Target: | CD28 |
| Reactivity: | Human |
| Method Type: | Sandwich ELISA |
| Application: | ELISA |

| Product Details | |
|--------------------|---|
| Purpose: | Custom Human CD28 ELISA Kit. |
| Sample Type: | Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Lysate |
| Analytical Method: | Quantitative |
| Detection Method: | Colorimetric |
| Specificity: | The antibody pair provided in this kit recognizes Human CD28 |
| Characteristics: | Strip plates and additional reagents allow for use in multiple experiments Quantitative protein detection Establishes normal range The best products for confirmation of antibody array data |
| Components: | Pre-Coated 96-well Strip Microplate Wash Buffer Stop Solution Assay Diluent(s) Lyophilized Standard Biotinylated Detection Antibody |

Product Details

- Streptavidin-Conjugated HRP
- TMB One-Step Substrate

Material not included:

- · Distilled or deionized water
- Precision pipettes to deliver 2 µL to 1 µL volumes
- Adjustable 1-25 µL pipettes for reagent preparation
- 100 µL and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- Log-log graph paper or computer and software for ELISA data analysis

Target Details

| Target: | CD28 |
|-------------------|--|
| Alternative Name: | CD28 (CD28 Products) |
| Gene ID: | 940 |
| UniProt: | P10747 |
| Pathways: | TCR Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of |
| | Molecular Mediator of Immune Response |

Application Details

| Sample Volume: | 100 μL |
|----------------|---|
| Plate: | Pre-coated |
| Protocol: | 1. Prepare all reagents, samples and standards as instructed in the manual. |
| | 2. Add 100 μL of standard or sample to each well. |
| | 3. Incubate 2.5 h at RT or O/N at 4 °C. |
| | 4. Add 100 μL of prepared biotin antibody to each well. |
| | 5. Incubate 1 h at RT. |
| | 6. Add 100 μL of prepared Streptavidin solution to each well. |
| | 7. Incubate 45 min at RT. |
| | 8. Add 100 μL of TMB One-Step Substrate Reagent to each well. |
| | 9. Incubate 30 min at RT. |
| | 10. Add 50 μL of Stop Solution to each well. |
| | 11. Read at 450 nm immediately. |
| | 11. Read at 450 nm immediately. |

Restrictions: For Research Use only

Handling

| Storage: | -20 °C |
|------------------|--|
| Storage Comment: | The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it is recommended to store at -80°C. |
| Expiry Date: | 6 months |