

# Datasheet for ABIN4885345

# **TFPI2 ELISA Kit**



### Overview

Quantity:	96 tests
Target:	TFPI2
Reactivity:	Human
Method Type:	Sandwich ELISA
Application:	ELISA

#### Product Details

Purpose:	Custom Human TFPI-2 SpeedELISA Kit.
Fulpose.	Custom Human 11 F1-2 SpeedLLISA Nt.
Brand:	SpeedELISA
Sample Type:	Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Lysate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Characteristics:	Compressed workflow with only one wash step and three incubation steps
	<ul> <li>3 hours total processing time</li> </ul>
	Strip plates and additional reagents allow for use in multiple experiments
	Quantitative protein detection
	Establishes normal range
	The best products for confirmation of antibody array data
Components:	Streptavidin-Coated 96-well Strip Microplate
	Wash Buffer
	Stop Solution
	Assay Diluent(s)

### **Product Details**

- · Lyophilized Standard
- · Biotinylated Capture Antibody
- · HRP Conjugated Detection Antibody
- TMB One-Step Substrate

#### Material not included:

- · Distilled or deionized water
- Precision pipettes to deliver 2  $\mu$ L to 1  $\mu$ L volumes
- Adjustable 1-25 µL pipettes for reagent preparation
- 100 µL and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- · Absorbent paper

TFPI2

- Microplate reader capable of measuring absorbance at 450nm
- · Log-log graph paper or computer and software for ELISA data analysis

### **Target Details**

Target:

11112
TFPI-2 (TFPI2 Products)
7980
P48307
100 μL
3 h
Pre-coated
1. Prepare all reagents, samples and standards as instructed in the manual.
2. Add 70 µL of the Antibody Cocktail to each well.
3. Incubate 10 minutes at RT.
4. Add 50 μL of sample or standard to each well.
5. Incubate 2 h at RT.
6. Wash Wells
7. Add 100 µL of TMB One-Step Substrate Reagent to each well.
8. Incubate 30 min at RT.
9. Add 50 µL of Stop Solution to each well.
10. Read at 450 nm immediately.
For Research Use only

# Handling

Storage:	-20 °C
Storage Comment:	May be stored for up to 6 months at 2° to 8°C from the date of shipment. Opened Microplate
	Wells or reagents may be store for up to 1 month at 2° to 8°C. Return unused wells to the
	pouch containing desiccant pack, reseal along entire edge. Reconstituted standard can be
	stored at -80°C for up to 1 week.
	Note: the kit can be used within one year if the whole kit is stored at -20°C. Avoid repeated
	freeze-thaw cycles.
Expiry Date:	6 months