antibodies -online.com





Datasheet for ABIN4885954

elF4EBP1 ELISA Kit



Overview

Quantity:	96 tests
Target:	elF4EBP1 (EIF4EBP1)
Binding Specificity:	pThr36, total
Reactivity:	Mouse, Human
Method Type:	Sandwich ELISA
Application:	ELISA
Product Details	
Purpose:	Human/Mouse Phospho-4E-BP1 (T36) and Total 4E-BP1 ELISA Kit. This assay semi-quantitatively measures phophorylated 4E-BP1 (Thr36) and Total 4E-BP1 in lysate samples.
Sample Type:	Cell Lysate, Tissue Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	The antibody pair provided in this kit recognizes human and mouse 4E-BP1 phosphorylated at site Threonine-36 and total 4E-BP1.
Characteristics:	 Simultaneously measure Phosphorylated protein and pan protein in one experiment (for normalization purpose) Screen numerous different cell lysates without performing a Western Blot analysis Minimal hands-on time, convenient, and non-radioactive material
Components:	Pre-Coated 96-well Strip MicroplateWash Buffer

Product Details

- · Anti-Phospho Antibody
- · Anti-Pan Antibody
- HRP-Conjugated Secondary Antibody
- · Streptavidin-Conjugated HRP
- · Assay Diluent
- · TMB One-Step Substrate
- · Stop Solution
- · Lysis Buffer
- · Positive Control Sample

Material not included:

- Distilled or deionized water
- · 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- · Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 µL to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- · Benchtop rocker or shaker
- · Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target:	elF4EBP1 (ElF4EBP1)
Alternative Name:	4E-BP1 (EIF4EBP1 Products)
Gene ID:	1978
UniProt:	Q13541
Pathways:	MAPK Signaling, PI3K-Akt Signaling, RTK Signaling, AMPK Signaling, Regulation of Cell Size, BCR Signaling

Application Details

Sample Volume:	100 μL
Plate:	Pre-coated
Protocol:	1. Prepare all reagents and samples as instructed in the manual.
	2. Add 100 μL of sample or positive control to each well.
	3. Incubate 2.5 h at RT or O/N at 4 °C.
	4. Add 100 μL of prepared primary antibody to each well.
	5. Incubate 1 h at RT.
	6. Add 100 μL of prepared 1X HRP-Streptavidin to each well.

Application Details

Expiry Date:

6 months

7. Incubate 1 h at RT. 8. Add 100 µL of TMB One-Step Substrate Reagent to each well. 9. Incubate 30 min at RT. 10. Add 50 µL of Stop Solution to each well. 11. Read at 450 nm immediately. Restrictions: For Research Use only Handling -20 °C Storage: Storage Comment: Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.