antibodies -online.com







Acetyl-CoA Carboxylase alpha ELISA Kit



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	96 tests	
Target:	Acetyl-CoA Carboxylase alpha (ACACA)	
Binding Specificity:	pSer79, total	
Reactivity:	Human, Rat	
Method Type:	Sandwich ELISA	
Application:	ELISA	
Product Details		
Purpose:	Human/Rat Phospho-ACC1 (S79) and Total ACC1 ELISA Kit. This assay semi-quantitatively	
	measures phophorylated ACC1 (Ser79) and Total ACC1 in lysate samples.	
Sample Type:	Cell Lysate, Tissue Lysate	
Analytical Method:	Semi-Quantitative	
Detection Method:	Colorimetric	
Specificity:	The antibody pair provided in this kit recognizes human, and rat ACC1 phosphorylated at site	
	Serine-79 and Total ACC1.	
Characteristics:	 Simultaneously measure Phosphorylated protein and pan protein in one experiment (for normalization purpose) 	
	Screen numerous different cell lysates without performing a Western Blot analysis	
	Minimal hands-on time, convenient, and non-radioactive material	
Components:	Pre-Coated 96-well Strip Microplate	
	Wash Buffer	

- · Anti-Phospho Antibody
- · Anti-Pan Antibody
- · HRP-Conjugated Secondary Antibody
- · Streptavidin-Conjugated HRP
- · Assay Diluent
- · TMB One-Step Substrate
- · Stop Solution
- · Lysis Buffer
- · Positive Control Sample

Material not included:

- · Distilled or deionized water
- · 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- · Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 µL to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- · Benchtop rocker or shaker
- · Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target:	Acetyl-CoA Carboxylase alpha (ACACA)	
Alternative Name:	ACC1 (ACACA Products)	
Gene ID:	31	
UniProt:	Q13085	
Pathways:	AMPK Signaling, Ribonucleoside Biosynthetic Process, Warburg Effect	

Application Details

, application betails	
Sample Volume:	100 μL
Plate:	Pre-coated
Protocol:	1. Prepare all reagents and samples as instructed in the manual.
	2. Add 100 µL of sample or positive control to each well.
	3. Incubate 2.5 h at RT or O/N at 4 °C.
	4. Add 100 μL of prepared primary antibody to each well.
	5. Incubate 1 h at RT.
	6. Add 100 µL of prepared 1X HRP-Streptavidin to each well.
	7. Incubate 1 h at RT.
	8. Add 100 µL of TMB One-Step Substrate Reagent to each well.

Application Details

	9. Incubate 30 min at RT.
	10. Add 50 μL of Stop Solution to each well.
	11. Read at 450 nm immediately.
Restrictions:	For Research Use only
Handling	
Storage:	-20 °C
Storage Comment:	Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of
	shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-
	Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell
	Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return
	unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20
	°C. Reconstituted Positive Control (Item K) should be stored at -70 °C.
Expiry Date:	6 months