Datasheet for ABIN4886014

c-FOS ELISA Kit

Go to Product page

Overview

Quantity:	96 tests
Target:	c-FOS (c-Fos)
Binding Specificity:	pThr232, total
Reactivity:	Human
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details

Purpose:	Human Phospho-C-Fos (Thr232) ELISA Kit. This assay semi-quantitatively measures phophorylated C-Fos (Thr232) in lysate samples.
Sample Type:	Cell Culture Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA kit recognizes Human C-Fos phosphorylated at site Threonine-232.
Characteristics:	• Rapidly measure phosphorylated protein in lysates • Screen numerous different cell lysates without performing a Western Blot analysis • Minimal hands-on time, convenient, and non-radioactive material
Components:	• Pre-Coated 96-well Strip Microplate • Wash Buffer • Anti-Phospho Antibody • HRP-Conjugated Secondary Antibody

Copyright antibodies-online. All rights reserved.

Product Details

	• Assay Diluent • TMB One-Step Substrate • Stop Solution • Lysis Buffer • Positive Control Sample
Material not included:	• Distilled or deionized water • 100 mL and 1 liter graduated cylinders • Tubes to prepare sample dilutions • Protease and Phosphatase inhibitors • Precision pipettes to deliver 2 µL to 1 mL volumes • Adjustable 1-25 mL pipettes for reagent preparation • Benchtop rocker or shaker • Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target:	c-FOS (c-Fos)
Alternative Name:	C-Fos (c-Fos Products)
Gene ID:	2353
UniProt:	P01100
Pathways:	S100 Proteins

Application Details

Sample Volume:	100 µL
Plate:	Pre-coated
Protocol:	1. Prepare all reagents and samples as instructed in the manual. 2. Add 100 µL of sample or positive control to each well. 3. Incubate 2.5 h at RT or O/N at 4 °C. 4. Add 100 µL of prepared primary antibody to each well. 5. Incubate 1 h at RT. 6. Add 100 µL of prepared 1X HRP-Streptavidin to each well. 7. Incubate 1 h at RT. 8. Add 100 µL of TMB One-Step Substrate Reagent to each well. 9. Incubate 30 min at RT. 10. Add 50 µL of Stop Solution to each well. 11. Read at 450 nm immediately.

Copyright antibodies-online. All rights reserved.

Application Details

Restrictions:	For Research Use only

Handling

Storage:	-20 °C
Storage Comment:	Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.
Expiry Date:	6 months

Copyright antibodies-online. All rights reserved.